

de Azevedo JR, de Araujo LO, da Silva WS, de Azevedo RP. A carbohydrate-restrictive strategy is safer and as efficient as intensive insulin therapy in critically ill patients. *J Crit Care*. 2009 Feb 12. [Epub ahead of print]

Doley RP, Yadav TD, Wig JD, Kochhar R, Singh G, Bharathy KG, Kudari A, Gupta R, Gupta V, Poornachandra KS, Dutta U, Vaishnavi C. Enteral nutrition in severe acute pancreatitis. *JOP*. 2009 Mar 9;10(2):157-62.

Hsu CW, Sun SF, Lin SL, Kang SP, Chu KA, Lin CH, Huang HH. Duodenal versus gastric feeding in medical intensive care unit patients: a prospective, randomized, clinical study. *Crit Care Med*. 2009 Jun;37(6):1866-72.

Kuhls DA, Rathmacher JA, Musngi MD, Frisch DA, Nielson J, Barber A, MacIntyre AD, Coates JE, Fildes JJ. Beta-hydroxy-beta-methylbutyrate supplementation in critically ill trauma patients. *J Trauma*. 2007 Jan;62(1):125-31; discussion 131-2.

Moses V, Mahendri NV, John G, Peter JV, Ganesh. Early hypocaloric enteral nutritional supplementation in acute organophosphate poisoning--a prospective randomized trial. *Clin Toxicol (Phila)*. 2009 May;47(5):419-24.

Savioli M, Cugno M, Polli F, Taccone P, Bellani G, Spanu P, Pesenti A, Iapichino G, Gattinoni L. Tight glycemic control may favor fibrinolysis in patients with sepsis. *Crit Care Med*. 2009 Feb;37(2):424-31.

White H, Sosnowski K, Tran K, Reeves A, Jones M. A randomised controlled comparison of early post-pyloric versus early gastric feeding to meet nutritional targets in ventilated intensive care patients. *Crit Care*. 2009;13(6):R187. Epub 2009 Nov 25.

Zhihui T, Wenkui Y, Weiqin L, Zhiming W, Xianghong Y, Ning L, Jieshou L. A randomised clinical trial of transnasal endoscopy versus fluoroscopy for the placement of nasojejunal feeding tubes in patients with severe acute pancreatitis. *Postgrad Med J*. 2009 Feb;85(1000):59-63.