




Brought to you by www.criticalcarenutrition.com and your ICU Dietitian

Consequences of the REDOXS and MetaPlus trials: the end of an era of glutamine and antioxidant supplementation for critically ill patients?

Recently, two large-scale, multicenter randomized trials studying the effect of supplemental glutamine and antioxidants in critically ill ventilated patients have been published^{1,2}. Together, the results of these 2 studies challenge current guidelines and recommendations for these special nutrients.

The REDOXS Trial was a factorial 2x2 randomized trial conducted in 40 intensive care units (ICUs) in North America and Europe¹. A total of 1223 mechanically ventilated adult patients with multi-organ failure were randomized to receive high doses of glutamine, antioxidants, both, or placebo administered separate from artificial nutrition. Contrary to the study hypothesis, the primary analysis demonstrated no clinical benefit of these nutritional interventions and identified a trend toward increased mortality at 28 days (32.4% vs. 27.2%; adjusted odds ratio, 1.28; 95% confidence interval [CI], 1.00 to 1.64; P = 0.049) and a significant increase in hospital and 6 month mortality among patients who received glutamine as compared with those who did not receive glutamine. Overall, there was no effect of antioxidants on 28-day mortality (30.8%, vs. 28.8%; adjusted odds ratio, 1.09; 95% CI, 0.86 to 1.40; P = 0.48). In a *post hoc* subgroup analysis, both glutamine and antioxidants appeared most harmful in patients with baseline renal dysfunction. No subgroups suggested reduced mortality with supplements³. The adverse effects of glutamine observed in the REDOXS trial were partially attributed to the high dose of study supplements provided to these patients (intravenous glutamine supplementation (0.35 g/kg/day parenterally provided as 0.50 g/kg/day of the dipeptide alanyl-glutamine [Dipeptiven®]) and an additional 30 g/day of glutamine enterally, provided as 42.5 g alanyl-glutamine and glycine-glutamine dipeptides)




Glutamine C₅H₁₀N₂O₃

The MetaPlus trial was conducted from February 2010 through April 2012 including a 6-month follow-up period in 14 ICUs in the Netherlands, Germany, France, and Belgium. A total of 301 adult patients who were expected to be ventilated and to require enteral nutrition (EN) for more than 72 hours were randomized to the intervention feed (High-protein EN enriched with 30 grams of glutamine, extra antioxidants including an additional 285 mcg of selenium and an additional 7.5 grams of fish oils (per 1500 ml), n = 152) or control (isocaloric standard high-protein EN (Nutrison Advanced Protison, NV Nutricia, Zoetermeer), n = 149) group initiated within 48 hours of ICU admission and continued during the ICU stay for a maximum of 28 days and included in an intention-to-treat analysis, performed for the total population as well as predefined medical, surgical, and trauma subpopulations.

There were no statistically significant differences in the primary end point, incidence of new infections according to the Centers for Disease Control and Prevention (CDC) definitions between the groups: 53% (95%CI, 44%-61%) in the enriched group vs 52% (95%CI, 44%-61%) in the control group (P = .96). Secondary end points included mortality, Sequential Organ Failure Assessment (SOFA) scores, mechanical ventilation duration, ICU and hospital lengths of stay, and subtypes of infections according to CDC definitions. No statistically significant differences were observed in other end points, except for a higher 6-month mortality rate in the medical subgroup: 54% (95%CI, 40%-67%) in the enriched group vs 35% (95%CI, 22%-49%) in the control group (P = .04), with a hazard ratio of 1.57 (95%CI, 1.03-2.39; P = .04) for 6-month mortality adjusted for age and Acute Physiology and Chronic Health Evaluation II score. The 6 month survival curves in this study in fact mirrored almost exactly the 6 month survival curves from the REDOXS study (see Figure 1).

In contrast, there have been 2 recent meta-analyses published that continue to show a positive treatment effect of supplemental glutamine and antioxidants^{4,5}. Wischmeyer and colleagues aggregated statistically 26 RCTs of IV glutamine supplementation involving 2484 patients. Trials of EN glutamine including the REDOXS and MetaPlus trial were not included in this review. They observed that IV glutamine supplementation was associated with a trend towards a reduction of overall mortality (RR 0.88, 95% CI 0.75, 1.03, P = .10) and a significant reduction in hospital mortality (RR 0.68, 95% CI 0.51, 0.90, P = .008). In addition, IV glutamine was associated with a strong trend towards a reduction in infectious complications (RR 0.86, 95% CI 0.73, 1.02, P = .09) and ICU length of stay (LOS) (WMD - 1.91, (95% CI -4.10, 0.28, P = .09) and significant reduction in hospital LOS (WMD -2.56, 95% CI -4.71, -0.42, P = .02). However, because of a concern about 'single-center' bias⁶, these investigators went on to show that only the single center trials demonstrated a significant effect of glutamine on overall and hospital mortality and infectious outcomes with no beneficial effect observed in the multicenter trials. In another recent glutamine meta-analysis, investigators compared the treatment effect of trials conducted over the last 2 decades and demonstrated that only trials done before 2003 manifested a positive signal while more recent trials failed to demonstrate any positive treatment effect⁷. Hence, it would appear that only old, small, single center trials of IV glutamine, when meta-analyzed showed a positive treatment effect.

Figure 1. 6-month probabilities of survival in the REDOXS and MetaPlus studies



With respect to antioxidants, Manzaneres and colleagues recently performed a meta-analysis of 21 RCTs of antioxidant supplementation in critically ill patients that included the results of the REDOXS study. When the results of these studies were statistically aggregated, combined antioxidants were associated with a significant reduction in mortality (risk ratio [RR]= 0.82, 95% confidence interval [CI] 0.72-0.93, P = .002); a significant reduction in duration of mechanical ventilation (weighed mean difference in days = -0.67, 95% CI -1.22,-0.13, P = .02); a trend towards a reduction in infections (RR= 0.88, 95% CI 0.76,1.02, P = .08); and no overall effect on ICU or hospital LOS. With few exceptions, most RCTs included in this systematic review were relatively small studies with a number of patients lower than 100, and thus inadequate to detect clinically important treatment effect of combined antioxidants on mortality. The signal emerges only after statistically aggregating these smaller trials.

How do we put this all together?

The signal of harm from these substrates comes from 2 large-scale, internally valid, multicenter trials that used both high dose IV and EN nutrients and a trial that used low dose EN nutrients only. In contrast, the signal of benefit seems to come from old, small, single-center trials. Small single center trials would be considered inadequate to guarantee safety as they are grossly underpowered to detect harm. These small trials, combined in meta-analyses, formed the basis of past clinical practice guidelines and now that larger scale trials with adequate power to evaluate mortality outcomes are conducted, these guidelines need to be updated. Given that our first dictum in medicine is to do no harm, we cannot be confident that supplemental glutamine and antioxidants are safe, whether provided enterally or parenterally, whether high or low dose. Until more data are available, we recommend that glutamine and antioxidants not be routinely administered to mechanically ventilated critically ill patients.

Key messages

1. There are now 2 studies that suggest that glutamine and antioxidants are harmful in critically ill patient populations.
2. Evidence in support of glutamine and antioxidants comes from old, single-centered RCTs and individually are inconclusive. The positive signal is only observed in meta-analysis of these RCTs which has not been confirmed in recent, large-scale, multi-center trials.
3. More research on the safety and efficacy of glutamine and antioxidants is needed before treatment recommendations can be considered.

References

1. Heyland D, Muscedere J, Wischmeyer PE, Cook D, Jones G, Albert M, Elke G, Berger MM, Day AG; Canadian Critical Care Trials Group. A randomized trial of glutamine and antioxidants in critically ill patients. *N Engl J Med*. 2013 Apr 18;368(16):1489-97.
2. van Zanten AR, Sztark F, Kaisers UX, Zielmann S, Felbinger TW, Sablotzki AR, De Waele JJ, Timsit JF, Honing ML, Keh D, Vincent JL, Zazzo JF, Fijn HB, Petit L, Preiser JC, van Horssen PJ, Hofman Z. High-protein enteral nutrition enriched with immune-modulating nutrients vs standard high-protein enteral nutrition and nosocomial infections in the ICU: a randomized clinical trial. *JAMA*. 2014 Aug 6;312(5):514-24.
3. Heyland DK, Elke G, Cook D, Berger MM, Wischmeyer PE, Albert M, Muscedere J, Jones G, Day AG; on behalf of the Canadian Critical Care Trials Group. Glutamine and Antioxidants in the Critically Ill Patient: A Post Hoc Analysis of a Large-Scale Randomized Trial. *JPEN*. 2014 May 5. [Epub ahead of print].
4. Wischmeyer PE, Dhaliwal R, McCall M, Ziegler TR, Heyland DK. Parenteral glutamine supplementation in critical illness: a systematic review. *Crit Care*. 2014 Apr 18;18(2):R76. [Epub ahead of print].
5. Manzaneres W, Dhaliwal R, Jiang X, Murch L, Heyland DK. Antioxidants micronutrients in the critically ill: a systematic review and meta-analysis. *Critical Care* 2012;16(2):R66.
6. Bellomo R, Warrillow SJ, Reade MC. Why we should be wary of single-center trials. *Crit Care Med*. 2009 Dec;37(12):3114-9.
7. Fadda V, Maratea D, Trippoli S, Messori A. Temporal trend of short-term mortality in severely ill patients receiving parenteral glutamine supplementation. *Clin Nutr*. 2013 Jun;32(3):492-3.