

9.2 Composition of Parenteral Nutrition: Type of lipids

Question: Does the type of lipids in parenteral nutrition affect outcomes in the critically ill adult patient?

Summary of evidence: A total of 21 level 2 studies (Nijveldt 1998, Lindgren 2001, Garnacho-Montero 2002, Garcia de Lorenzo 2005, Iovinelli 2007, Guo 2008, Wang 2009, Qu 2009, Barbosa 2010, Sabater 2011, Khor 2011, Zhao 2011, Pontes-Arruda 2012, Burkhart 2013, Gultekin 2014, Hall 2014, Wang 2014, Chen 2017a, Chen 2017b, Donaghue 2019 and Singer 2021) and 5 level 1 studies (Grecu 2003, Friesecke 2008, Gupta 2011, Umperrez 2012 & Grau Carmona 2014) were reviewed. For most of the studies, the focus of the investigation was on surrogate endpoints, but the studies were still included because they did report clinical outcomes such as mortality or infections.

Twenty-one trials using parenteral nutrition compared varying strategies of reducing omega-6 fatty acids from soybean oil (SO), and 5 trials used FO as a stand-alone supplement (Gupta 2011, Khor 2011, Zhao 2011, Burkhart 2014 and Hall 2015)

Omega-6 reducing lipid emulsions: SO/MCT or SO/OO

Four trials compared mixtures of SO (long-chain triglycerides [LCTs]) plus medium chain triglycerides (MCT) (SO/MCT), to a SO (LCT) emulsion (Nijveldt 1998, Lindgren 2001, Garnacho-Montero 2002 and Iovinelli 2007). Additionally, 3 studies compared a mixture of soybean and olive oil (SO/OO, Clinoleic) to a SO/MCT mixture or SO (Garcia de-Lorenzo 2005, Umperrez 2012 & Pontes-Arruda 2012). The trial by Pontes-Arruda include 3 groups: 1) multichambered bag (MCB) with OO/SO, 2) compounded PN with OO/SO and 3) compounded SO. In the meta-analysis, we have only included group 2 and 3 so as to evaluate the effect of OO and not effect of the MCB. The trial by Huschak was excluded from the meta-analysis because the PN strategies also varied by lipid and glucose composition

FO-containing lipid emulsions

The goal of FO-containing lipid emulsions was not only to lower amount the amount linoleic acid but to create a more favourable omega-3-to-omega-6 fatty acid ratio by adding a FO supplement to PN or either adding a FO containing lipid component to MCT/LCT (Lipoplus/Lipidem) or to blend of MCT/LCT and OO (SMOFlipid).¹

¹ it is important to know that the fish oil enriched component of the two commercially available products Lipoplus and SMOFlipid comply with different European pharmacopeia (EP) monographs. Lipoplus complies with EP monograph 1352 "Omega-3-acid triglycerides" which is defined as a synthesized mixture of mono-, di- and triesters of omega-3 acids. Sum of the contents of the omega-3 acids EPA and DHA expressed as triglycerides is not less than 45%. SMOFlipid complies with EP monograph 1912 "Fish oil, rich in omega-3 acids" which describes purified fish oil rich in omega-3 fatty acids. The monograph specifies that the content of EPA is not less than 13% and a content of DHA not less than 9%. Lipoplus contains 10% omega-3 acid triglycerides and SMOFlipid contains 15% fish oil, rich in omega-3 acids. Although omega-3 fatty acids are derived from natural sources (leading to slight variations in concentration), the actual concentrations of EPA and DHA in SMOFlipid and Lipoplus are comparable according to existing literature. Both products provide approximately the same omega-3-to-omega-6 (ratio of 1:2.5 to 1:2.6).

Seven trials of patients receiving parenteral nutrition compared lipid emulsions supplemented with fish oil (FO) (i.v. FO supplement Omegaven) to SO, MCT/SO or OO/SO (Greco 2003, Friesecke 2008, Wang 2009, Gultekin 2014, Wang 2014, Chen 2017a, and Chen 2017b). The added FO emulsion was not specified in two trials (Guo 2009 and Qu 2009). Four trials compared a ready-to-use mixture of SO, MCT and FO² (SO/MCT/FO, Lipoplus) to SO/MCT (Barbosa 2010, Grau-Carmona 2014, Singer 2021) or SO (Sabater 2011), and 1 study compared another ready-to-use mixture of SO, MCT, olive oil (OO) and FO³ (SO/MCT/OO/FO, SMOFlipid) to LCT (Donoghue 2019).

Singer 2021 used an EN product enriched with fish oils in the intervention group, which was different than all other trials. The dose from parenteral fish oils was initially low (<0.05 g/kg/d on d1-5, >0.05 g/kg/d from d6) and most of the fish oils were derived from the enteral product so the effect of this trial was evaluated in a *sensitivity analysis* where it was excluded (See Table 2 on Fish Oil Dosing).

One study that compared an outdated long chain triglyceride (LCT) emulsion to another form of LCT (Kari 1998) was removed in the 2013 CPGs as it did not involve a soybean oil reducing strategy. The Wang 2008 study was replaced by a later version of the study by the same authors that had more patients i.e., Wang 2009. All of the studies had a goal of reducing the amount of omega-6 fatty acids in the setting of PN use.

Another 5 trials compared stand alone fish oil enriched IV lipid emulsions in patients fed as per usual clinical routine where the control patients received standard of care (Gupta 2011, Khor 2011, Zhao 2011, Burkhart 2013 and Hall 2014). These trials were analyzed separately.

Subgroup Analyses

We compared the effect of SO/MCT or SO/OO vs SO and FO-containing PN vs SO or SO/MCT or SO/OO. In addition, as Omegaven (10% lipid emulsion from pure fish oils) was used in 12 trials (7 were PN-fed trials and 5 were in the context of a stand-alone supplement), we performed a comparison of the trials of Omegaven vs. non-Omegaven fish oil containing strategies.

² FO component as Omega-3 Acid Triglycerides according to Pharm Eur Monograph 1352

³ FO component as Fish oil, Rich in EPA and DHA according to Pharm Eur Monograph 1912

Mortality:

Overall omega-6 fatty acid reducing strategy: When all the trials that used an omega-6 fatty acid sparing strategy, excluding those in which the FO supplements were given as a stand-alone intervention, were aggregated, the use of a lower omega-6 fatty acid strategy was not associated with a significant reduction in overall mortality (RR 0.91, 95% CI 0.76, 1.10, $p=0.34$, test for heterogeneity $I^2 = 0\%$; figure 1.1). However, a trend towards reduction in 28-day mortality was observed (RR 0.79, 95% CI 0.61, 1.02, $p=0.07$, $I^2 = 0\%$; figure 2.1)

When sensitivity analysis was done without the Singer 2021 trial, a similar results was observed for overall mortality (RR 0.91, 95% CI 0.75, 1.11, $p=0.36$, test for heterogeneity $I^2 = 0\%$; figure 1.2), and 28-day mortality (RR 0.77, 95% CI 0.59, 1.02, $p=0.07$, $I^2 = 0\%$; figure 2.2)

In subgroup analyses, neither SO/MCT or SO/OO vs OO (RR 0.89, 95% CI 0.65, 1.23, $p=0.50$, test for heterogeneity $I^2 = 0\%$; figure 1.3) nor FO-containing PN vs SO or SO/MCT or SO/OO (RR 0.92, 95% CI 0.72, 1.16, $p=0.47$, test for heterogeneity $I^2 = 4\%$; figure 1.3) had an effect on overall mortality. The test for subgroup differences were non-significant ($p=0.91$).

A trend towards reduction of 28-day mortality was found for FO-containing PN vs SO or SO/MCT or SO/OO (RR 0.74, 95% CI 0.54, 1.01, $p=0.06$, $I^2 = 0\%$; figure 2.3). The test for subgroup differences were non-significant ($p=0.50$).

SO/MCT vs. SO: A meta-analysis of the studies of SO/MCT vs. SO showed no difference in overall mortality between the groups (RR 0.90, 95% CI 0.55, 1.48, $p=0.68$, heterogeneity $I^2=0\%$; figure 1.1). No studies reported 28-day mortality (figure 2.1)

SO/OO vs SO/MCT or SO: We observed no difference in overall mortality (RR 0.89, 95% CI 0.59, 1.34, $p = 0.57$, heterogeneity $I^2=0\%$; figure 1.1) and 28-day mortality (RR 0.91, 95% CI 0.56, 1.47, $p = 0.69$, figure 2.1) between the groups receiving the OO containing emulsions compared to SO/MCT or SO

FO containing PN vs. SO or SO/MCT or SO/OO: With respect to studies of FO enriched PN vs. SO or SO/MCT or SO/OO, no significant reduction in overall mortality was observed (RR 0.92, 95% CI 0.72, 1.16, $p = 0.47$, test for heterogeneity $I^2 =4\%$; figure 1.1). In the sensitivity analyses without the Singer 2021 trial, the result was similar (RR 0.90, 95% CI 0.69, 1.18, $p = 0.46$, test for heterogeneity $I^2 =12\%$; figure 1.2). However, a trend towards lower 28-day mortality was observed in the main (RR 0.74, 95% CI 0.54, 1.01, $p=0.06$, $I^2 =0\%$; figure 2.1 and sensitivity (RR 0.72, 95% CI 0.51, 1.01, $p=0.05$, $I^2 =0\%$; figure 2.2) analyses.

Omegaven vs. other oils:

FO containing PN: Omegaven was associated with a significant reduction in overall (RR 0.68, 95% CI 0.48, 0.95, $p=0.03$, test for heterogeneity $I^2 =0\%$; figure 1.4) and 28-day mortality (RR 0.63, 95% CI 0.41, 0.99, $p=0.05$, $I^2 =16\%$; figure 2.4) while other fish oil emulsions had no effect on overall (RR 1.19, 95% CI 0.87, 1.62, $p=0.27$, test for heterogeneity $I^2 =0\%$; figure 1.4) and 28-day (RR 0.91, 95% CI 0.56,

1.51, $p=0.74$, $I^2=0\%$; figure 2.4) mortality. The test for subgroup differences was significant for overall ($p=0.02$) but not 28-day ($p=0.28$) mortality..

In the sensitivity analyses that excluded the Singer 2021 trial, the effects of other fish oil emulsions on overall mortality were similar for both overall (RR 1.26, 95% CI 0.90, 1.77, $p=0.18$, test for heterogeneity $I^2=0\%$; figure 1.5) and 28-day mortality (RR 0.94, 95% CI 0.49, 1.83, $p=0.86$, $I^2=0\%$; figure 2.5), test for subgroup differences was significant in overall ($p=0.01$) but not 28-day mortality ($p=0.33$).

Stand-alone FO emulsion supplement vs. standard care: When these FO trials in which the control group received no lipids were aggregated, a trend towards significant reduction in overall mortality (RR 0.76, 95% CI 0.53, 1.10, $p=0.14$, test for heterogeneity $I^2=0\%$; figure 1.6), and a significant reduction of 28-day mortality (RR 0.60, 95% CI 0.36, 0.99, $p=0.04$, $I^2=0\%$; figure 2.6) were found. All aggregated trials used omegaven FO.

In studies that reported **ICU mortality**, no differences between groups were found in the overall and the subgroup analyses. None of the trial on stand-alone FO emulsion reported ICU mortality (figure 3.1 to 3.4)

In studies that reported **hospital mortality**, no differences no differences between groups were found in the overall and the subgroup analyses (figure 3.1-3.3). A trend towards reduced hospital mortality was found when the 2 trials of stand-alone FO supplement were aggregated (RR 0.59, 95% CI 0.34, 1.04, $p=0.07$, $I^2=0\%$; figure 4.4)

Infections:

Overall omega-6 fatty acid reducing strategy: When all 7 studies that used an omega-6 fatty acid sparing strategy were aggregated, the use of a lower omega-6 fatty acid emulsion had no effect on infections (RR 0.94, 95% CI 0.70, 1.26, $p = 0.68$, $I^2=32\%$; figure 5.1).

In subgroup analyses, **SO/OO vs SO/MCT or SO** were associated with a trend towards an increase in overall infections (RR 1.23, 95% CI 0.92, 1.63, $p=0.16$, $I^2 = 0\%$; figure 5.1) while **FO-containing PN** were associated with a significant reduction in infections (RR 0.65, 95% CI 0.44, 0.95, $p=0.03$, $I^2 = 0\%$; figure 5.1). There was a significant difference between the two subgroups ($p=0.008$).

Omegaven vs. other fish oils:

FO enriched PN: When the data from 3 studies of Omegaven were aggregated, there was no effect on overall infections (RR 0.77, 95% CI 0.44, 1.36, $p=0.37$, $I^2=0\%$; figure 5.2) while in one study a significant reduction in infections was seen in the group receiving fish oil emulsion other than Omegaven (Grau-Carmona 2014, RR 0.56, 95% CI 0.34, 0.94, $p=0.03$; figure 5.2). Test for subgroup differences was not significant ($p=0.43$).

FO vs. standard (no lipids): only one trial that used Omegaven was found, hence a subgroup comparison to non Omegaven studies was not possible (figure 5.2).

Stand-alone FO emulsion supplement vs. standard care: When examining the only trial of fish oils in which the control group received no IV soybean oil, no effect was seen on infections (Hall 2014, RR 0.60, 95% CI 0.16-2.29; $p=0.45$; no figure shown).

Hospital LOS:

Overall omega-6 fatty acid reducing strategy: When the 6 studies that used an omega-6 fatty acid sparing strategy were aggregated, the use of a lower omega-6 fatty acid emulsion was associated with a significant reduction in hospital LOS (WMD -6.88, 95% CI -11.27, -2.49, $p=0.002$, test for heterogeneity $I^2=20\%$; figure 6.1).

In subgroup analyses, **SO/OO vs SO/MCT or SO** had no significant effect on hospital LOS (WMD -6.80, 95% CI -19.17, 5.57, $p=0.28$, test for heterogeneity $I^2= 0\%$; figure 3.2). A trend towards shorter hospital LOS was observed with **FO enriched PN vs. SO or SO/MCT or SO/OO** (WMD -5.93, 95% CI -13.13, 1.27, $p=0.11$, test for heterogeneity $I^2= 51\%$; figure 6.1). The test for subgroup differences was not significant ($p=0.90$).

Omegaven vs. other fish oils:

FO enriched PN: When the data from 2 studies of Omegaven were aggregated, there was no effect on hospital length of stay (WMD -5.75, 95% CI -14.61, 3.11; $p=0.20$, test for heterogeneity $I^2=61\%$; figure 6.2) and a similar lack of effect was seen in the aggregated data from the two studies of other fish oil emulsions (WMD -12.87, 95% CI -42.65, 16.91, $p=0.40$, test for heterogeneity $I^2=92\%$; figure 3.3). Test for subgroup differences was not significant ($p=0.65$).

FO vs. standard (no lipids): all three studies used Omegaven, hence a subgroup comparison to non Omegaven studies was not possible.

Stand-alone FO emulsion supplement vs. standard care: Fish oil emulsions had no effect on hospital LOS when compared to standard care (WMD = 0.78, 95% CI -2.89, 4.46, $p=0.68$, test for heterogeneity $I^2= 0\%$; figure 6.3).

ICU LOS

Overall omega-6 fatty acid reducing strategy: When all 12 studies that used an omega-6 fatty acid sparing strategy were aggregated, the use of a lower omega-6 fatty acid emulsion was associated with a trend towards a reduction in ICU LOS (WMD -1.94, 95% CI -4.41, 0.52, $p=0.12$, test for heterogeneity $I^2=83\%$; figure 7.1).

In subgroup analysis, SO/MCT or SO/OO vs SO had no effect on ICU LOS (WMD 1.74, 95% CI -2.17, 5.66, $p=0.38$, test for heterogeneity $I^2=50\%$; figure 7.2); however, FO-containing PN vs SO or SO/MCT or SO/OO significantly reduced ICU LOS (WMD -3.53, 95% CI -6.16 to -0.90, $p=0.009$, $I^2=82\%$; figure 7.2) There was a significant difference between the two subgroups ($p=0.03$).

SO/MCT vs SO: No difference in ICU LOS between the two groups (WMD 3.03, 95% CI -2.02, 8.07, $p=0.24$; figure 7.1).

SO/OO vs. SO/MCT or SO: When the data from the two studies of olive oil emulsions vs SO/MCT or SO were aggregated, olive oil emulsions had no effect on ICU length of stay (WMD -2.09, 95% CI -12.17, 8.00, $p=0.69$, test for heterogeneity $I^2=55\%$; figure 7.1).

FO containing PN vs. SO or MCT/SO or OO/SO: When the data from the eight studies of fish oil emulsions vs SO+ MCT or SO were aggregated, a significant reduction in ICU LOS was observed in the fish oil group (WMD -3.53, 95% CI -6.16 to -0.90, $p=0.009$, $I^2=82\%$; figure 7.1).

Omegaven vs. other fish oils:

FO enriched PN: When the data from 4 studies of Omegaven were aggregated, there was a trend towards reduction in ICU length of stay (WMD -3.15, 95% CI -6.89, 0.59, $p=0.10$, test for heterogeneity $I^2=79\%$; figure 7.3). Similarly, when the data were aggregated from the four studies of other fish oil emulsions, a trend towards reduction in ICU LOS was observed (WMD -3.81, 95% CI -7.96, 0.34, $p=0.07$, test for heterogeneity $I^2=75\%$; figure 7.3). Test for subgroup differences was not significant ($p=0.82$).

FO vs. standard (no lipids): all four studies used Omegaven, hence a subgroup comparison to non Omegaven studies was not possible.

Stand-alone FO emulsion supplement vs. standard care: When these trials were statistically aggregated, there was a trend towards reduced ICU LOS (WMD -1.38, 95% CI -4.11, 1.34, $p=0.32$; test for heterogeneity $I^2=52\%$; figure 7.4).

Ventilator days:

Overall omega-6 fatty acid reducing strategy: A trend towards shorter duration of ventilation was found with Omega-6 fatty acid sparing strategies (WMD -0.87, 95% CI -1.82, 0.07, $p=0.07$, test for heterogeneity $I^2=52%$; figure 8.1).

In subgroup analyses, SO/MCT vs SO/OO vs SO were associated with a significant reduction in ventilator days (WMD -3.26, 95% CI -5.32, -1.20, $p=0.002$, test for heterogeneity $I^2=0%$; figure 8.2) while FO-containing PN had no effect on ventilator days (WMD -0.31, 95% CI -1.07, 0.45, $p=0.42$, test for heterogeneity $I^2=39%$; figure 8.2). The test for subgroup differences were significant ($p=0.009$).

SO/MCT vs SO: Only one study comparing SO+MCT to SO reported duration of ventilation and a significant reduction duration of mechanical ventilation in the SO+MCT group was observed (RR -3.30, 95% CI -5.39, -1.21, $p=0.002$; figure 8.1).

SO/OO vs. SO/MCT or SO: Only one study reported this outcome, and the use of olive oil emulsions had no effect on the duration of mechanical ventilation (WMD -2.00, 95% CI -13.92, 9.92, $p=0.74$; figure 8.1).

FO containing PN vs. SO or SO/MCT or SO/OO: When the data from the six studies of fish oils were aggregated, no effect on duration of mechanical ventilation was observed (WMD -0.31, 95% CI -1.07, 0.45, $p=0.42$, test for heterogeneity $I^2=39%$; figure 8.1).

Omegaven vs. other fish oils:

FO enriched PN: When the data from 3 studies of Omegaven were aggregated, there was no effect on ventilator days (WMD -0.87, 95% CI -2.37, 0.63, $p=0.26$, test for heterogeneity $I^2=35%$; figure 8.3). Similar signals were seen when the three studies of other fish oil emulsions, were pooled (WMD 0.28, 95% CI -0.32, 0.88, $p=0.36$, test for heterogeneity $I^2=0%$; figure 8.3). Test for subgroup differences was not significant ($p=0.16$).

FO vs. standard (no lipids): both studies that reported on ventilator days used Omegaven, hence a subgroup comparison to non Omegaven studies was not possible.

Stand-alone FO emulsion supplement vs. standard care: When these two trials were statistically aggregated, no effect on ventilator days was observed (WMD 1.98, 95% CI -2.36, 6.31, $p=0.37$; test for heterogeneity $I^2=0%$; figure 8.4).

Other outcomes:

SO/MCT vs SO: A significant improvement in nutritional parameters (i.e., nitrogen balance, retinol binding protein, prealbumin) was observed in the groups receiving SO + MCT in some of the studies (Garnacho-Montero 2002, Lindgren 2001) and a significant reduction in the time of weaning was seen in one study (Iovinelli 2007).

SO/OO vs. SO/MCT or SO: No differences were found for multiple organ dysfunction score (Garcia-de-Lorenzo 2005) or other nutritional outcomes (Pontes-Arruda 2013, Umpierrez 2012)

FO containing PN vs. SO or SO/MCT or SO/OO: The use of FO was associated with a reduction in the need for surgery due to a subsequent septic episode when compared to LCT ($p=0.010$, Grecu 2003). Wang 2009 reported a reduction in the need for surgery for pancreatic necrosis in the group receiving FO, but this was not statistically different. There was a trend towards a reduction in catheter related blood stream infections in the group receiving fish oils ($p=0.10$, Friesecke 2008) and better gas exchange (Barbosa 2010). Singer 2021 reported a higher % patients tolerating EN alone at day 6 and higher catecholamine treatment free days in the omega 3 supplemented EN+PN group ($p=0.34$ and 0.05 respectively).

Conclusions: In critically ill patients,

- 1) Omega-6 FFA reducing strategies, also known as SO sparing strategies,
 - a. are not associated with a significant reduction in overall, ICU or hospital mortality, but a trend towards lower 28-day mortality was observed
 - b. are associated with a reduction in hospital LOS.
 - c. may be associated with a reduction in ICU LOS and duration of mechanical ventilation
 - d. have no effect on infections
- 2) SO/MCT emulsions, compared to SO, have no effect on mortality, infections, ICU or hospital length of stay in critically ill patients. There may be a significant reduction in duration of mechanical ventilation associated with SO/MCT but data points are too sparse to be sure (1 trial).
- 3) OO containing emulsions, compared to SO/MCT or SO,
 - a. have no effect on mortality or ICU/hospital LOS or duration of mechanical ventilation.
 - b. may be associated with increased infections
- 4) FO containing PN vs. SO or MCT/SO or OO/SO:
 - a. are not associated with a significant reduction in overall, ICU or hospital mortality but a trend towards lower 28-day mortality was observed
 - b. are associated with a significant reduction in infectious complications and ICU LOS.
 - c. may be associated with a reduction in duration of mechanical ventilation
 - d. have no effect on hospital LOS
- 5) Stand-alone FO emulsion supplement vs. standard care
 - a. is associated with a reduction of 28-day mortality and a trend towards reduction in overall and hospital mortality/
 - b. may be associated with a reduction in ICU LOS
 - c. have no effect on ICU mortality, infectious complications, duration of mechanical ventilation, or hospital LOS.
- 6) Compared to SO/MCT or SO/OO, FO-containing lipid emulsions
 - a. are associated with a significant reduction in infectious complications and ICU LOS
 - b. have no impact on mortality, hospital LOS and ventilator days.
- 7) Compared to non-Omegaven containing PN or IV nutritional strategies, Omegaven containing PN or IV strategies
 - a. May be associated with a significant reduction in mortality.

Level 1 study: if all of the following are fulfilled: concealed randomization, blinded outcome adjudication and an intention to treat analysis.

Level 2 study: if any one of the above characteristics is unfulfilled.

Table 1. Randomized trials evaluating type of lipids (PN) in critically ill patients

Study	Population	Methods (score)	Intervention	Mortality # (%)†		Infections # (%)‡	
(A) SO/MCT vs. SO							
1) Nijveldt 1998	ICU, septic surgical patients, trauma N=20	C.Random: not sure ITT: yes Blinding: double (9)	PN + Lipofundin (50% MCT+ 50% SO) vs. PN + Intralipid (100% SO)	MCT/SO Not specified 2/12 (17)	SO Not specified 1/8 (13)	MCT/SO NR	SO NR
2) Lindgren 2001	ICU patients, sepsis, multi-trauma N=20	C.Random: yes ITT: No Blinding: double (7)	PN + Structolipid (36% MCT/ 64%SO) vs. PN + Intralipid (100% SO)	MCT/SO Hospital 4/9 (44.4)	SO Hospital 5/11 (45.5)	NR	NR
3) Garnacho-Montero 2002	Surgical ICU Patients with peritonitis and abdominal sepsis N=72	C.Random: No ITT: Yes Blinding: no (4)	PN + Lipofundin (50% MCT + 50% SO) vs. PN with Intralipid (100% SO) Both groups received PN with 45 % Branched chain amino acids	MCT/SO ICU 8/35 (23) Hospital 11/35 (31)	SO ICU 11/37 (30) Hospital 13/37 (35)	MCT/SO NR	SO NR
4) Iovinelli 2007	Patients with COPD requiring ventilation N=14 (of 24)	C.Random: not sure ITT: no Blinding: no (4)	PN + Lipofundin (50% MCT + 50% SO) vs. 100% SO. In both received 50% of non-protein calories given as lipids	MCT/SO 15-d 2/7 (28.6)	SO 15-d 3/7 (42.9)	MCT/SO Catheter-related 1/12 (8)	SO Catheter-related 2/12 (17)
(B) SO/OO vs. SO/MCT or SO							
5) Garcia-de-Lorenzo 2005	Severe burn patients, burn severity index ≥ 7, TBSA > 30 % N=22	C.Random: not sure ITT: yes Blinding: double (10)	PN with ClinOleic 20% (80% OO, 20% SO, (63% ω9, 37% ω6= restricted linoleic acid {ω6} content) vs. Lipofundin (50% MCT+ 50% SO).	OO/SO ICU=Hosp=6-mo 4/11 (36)	MCT/SO ICU=Hosp=6-mo 3/11 (27.3)	OO/SO 6/11 (55)	MCT/S 6/11 (55)

6) Pontes-Arruda 2012	ICU pts requiring PN from 8 ICUs and 3 countries N=204	C.Random: yes ITT: yes Blinding: no (9)	PN with compounded ClinOleic 80% OO, 20% SO (n=103) vs compounded PN with a MCT/SO (n=101)	OO/SO ICU 19/103 (24) 28-day=Hosp* 24/103 (27)	MCTSO ICU 21/101 (21) 28-day=Hosp* 26/101 (26)	OO/SO 39/103 (38) All infections ICU acquired infections 28/103 (27) VAP/lower respiratory infections 9/103 (9)	MCT/SO 35/101 (35) ICU acquired infections 23/101 (23) ICU acquired infections 23/101 (23) VAP/lower respiratory infections 11/101 (11)
7) Umpierrez 2012	Medical surgical ICU pts post op (88% emergency surgeries) N=100	C.Random: yes ITT: yes Blinding: double (14)	PN with ClinOleic 20% (80%OO, 20% SO,, $\omega 6:\omega 3=9:1$) vs Intralipid (100% SO, $\omega 6:\omega 3=7:1$)	OO/SO Hospital 5/51 (10)	SO Hospital 8/49 (16)	OO/SO 29/51 (57) 7/51 (14)	SO 21/49 (43) Pneumonia 5/49 (10)
(C) FO containing PN vs. SO or SO/MCT or SO/OO							
8) Grecu 2003*	Patients with abdominal sepsis N=54 (15/54 in ICU)	C.Random: yes ITT: yes Blinding: double (12)	PN + Omegaven (10% fish oils) plus SOs vs. PN with SO	FO/SO ICU 2/28 (7)	SO ICU 3/26 (12)	FO/SO VAP 0/8	SO VAP 1/7 (14)
9) Friesecke 2008	Medical ICU patients N=166	C.Random: yes ITT: yes Blinding: double (10)	PN + Lipofundin (50% MCT + 50% SO) + Omegaven (10% fish oil) vs. Lipofundin MCT (50% MCT + 50% SO)	FO/MCT/SO 28 day 18/83 (22)	MCT/SO 28 day 22/82 (27)	FO/MCT/SO 10/83 (12)	MCT/SO 11/82 (13)
10) Guo 2008	Septic ICU patients with APACHE II > 12 N=80 (of 88)	C.Random: no ITT: no Blinding: no (4)	PN with 20% lipid emulsion with an added 100 ml of Omega-3 PUFAs (product not specified) vs PN with 20% lipid emulsion	FO 28 day 6/38	SO 28 day 8/42	NR	NR
11) Qu 2009	Severe sepsis patients N=40	C.Random: no ITT: no Blinding: no (5)	Routine PN + omega 3 FO emulsion (product not specified) at 1-2 ml/kg/d vs routine PN.	FO 28 day 4/20 (20)	Standard 28 day 2/20 (10)	NR	NR
12) Wang 2009	Severe acute pancreatitis patients in ICU N=56	C.Random: no ITT: yes Blinding: double (11)	PN + Omegaven (10% FO) plus Lipovenos (SO) ($\omega 3:\omega 6$ ratio was 1:4) vs. PN with Lipovenos (SO). Both received same amounts of lipids (1 gm/kg/day) Note: Lipovenos contains 20% MCT	FO ICU 0/28 (0)	SO ICU 2/28 (7)	FO 6/28 (21)	SO 9/28 (32)

13) Barbosa 2010	ICU patients with SIRS or sepsis requiring PN N=25	C.Random: yes ITT: yes Blinding: single (10)	PN + Lipolus (50% MCT, 40%SO, 10% FO) vs. Nutriflex LipidSpecial (50% MCT, 50% SO). Both received same amounts of lipids (~1 gm/kg/day)	FO/MCT/SO 5 day 2/13 (15) 28 day 4/13 (31)	MCT/SO 5 day 1/10 (10) 28 day 4/10 (40)	FO/MCT/SO NR	MCT/SO NR
14) Sabater 2011	ARDS pts requiring MV and PN	C.Random: unknown ITT: yes Blinding: unknown (9)	Group A: Lipoplus 20% B. Braun Medical (50% MCT, 40% LCT, 10% FO). Group B: Intralipid 20% Fresenius Kabi (100% LCT). The lipids emulsions were administered for 12h at 0.12 g/kg/h in both groups	FO/MCT/SO not specified 4/8 (50)	SO not specified 2/8 (25)	NR	NR
15) Grau Carmona 2014	Medical and surgical pts requiring TPN N=159 (of 175)	C.Random: yes ITT: no Blinding: double (8)	PN + Lipoplus (50% MCT, 40% SO, 10% FO vs PN + Lipofundin (50% MCT + 50% SO)	FO/MCT/SO ICU 26/81 (32.5) Hospital 32/81 (39.5) 6-month 34/81 (42)	MCT/SO ICU 16/78 (20.5) Hospital 22/78 (28.3) 6-month 24/78 (30.8)	FO/MCT/SO 17/81 (21)	MCT/SO 29/78 (37.2)
16) Gultekin 2014	ICU pts needing TPN N=32 (of 58)	C.Random: unknown ITT: other Blinding: double (4)	PN + 100ml/day Omegaven (10% FO) plus Clinoleic (80% OO, 20% SO) vs PN + Clinoleic. Both groups were prescribed IV lipids to provide 30-40% of total energy requirements.	FO/OO/SO Not specified 7/16 (44)	OO/SO Not specified 8/16 (50)	NR	NR
17) Wang 2014	Abdominal sepsis patients needing TPN for ≥5 days N=53	C.Random: unknown ITT: yes Blinding: unknown (6)	Routine PN + Omegaven 0.2 g/kg/day to replace part of routine PN lipid emulsion for 5 days vs Routine PN	NR	NR	NR	NR
18) Chen 2017a	ICU patients with SIRS N=78	C.Random: unknown ITT: yes Blinding: single (9)	PN containing 50g LCFA + 100 m/day containing 10g FO as Omegaven vs PN containing 50g LCFA. Both groups dosed at 20 kcal/kg for first 7 days, slowly increased to 30 kcal/kg afterwards	FO/SO 28 day 10/41 60 day 11/41	SO 28 day 15/37 60 day 18/37	NR	NR
19) Chen 2017b	Mechanically ventilated patients with systemic inflammation reaction syndrome and intestinal failure N=48	C.Random: no ITT: yes Blinding: double (9)	PN with 10 gms/day Omega-3 fatty acids (100 mls Omegaven) vs. standard PN (assumed to be LCT) Isocaloric (20 Kcal/kg/day in first 7 days)	FO 28 day 3/24 (12.5%)	SO 28 day 10/24 (41.7%) p =0.023	NR	NR

20) Donoghue 2019	Critically ill patients with SIRS, sepsis and/or ARDS N=68 (of 75)	C.Random: no ITT: no Blinding: double (7)	PN with SMOF lipid emulsion (30% SO, 30% MCT, 25% OO and 15% FO, $\omega 6:\omega 3=2.5:1$) vs. LCT Intralipid (100% SO $\omega 6:\omega 3=7:1$)	SMOF ICU 6/35 (18.4%)	SO ICU 5/33 (15.2%); p=0.71	NR	NR
21) Singer 2021	Mechanically ventilated patients, receiving <80% by EN alone N=95 (of 100)	C.Random: yes ITT: no Blinding: double (9)	Supplemental PN (Nutriflex Omega Special) with MCT/SO: Omega-3 PUFA of 4:5:1 (EPA+DHA: 1.8 g/ml) plus EN enriched with fish oils, borage oil & gamma linolenic acid (Oxepa) vs. Supplemental PN (Nutriflex lipid) with MCT/SO of 1:1 plus EN without fish oils (Pulmocare)	FO EN+PN 28 day 10/48 (20.8%) 90 day 15/48 (31.3%)	MCT/SO 28 day 11/47 (23.4%); p=0.81 90 day 19/47 (40.4%); p=0.52	NR	NR
(D) Stand-alone FO emulsion supplement vs. standard care							
22) Gupta 2011	ICU patients with suspected ARDS N=61	C.Random: yes ITT: yes Blinding: double (9)	EN (standard diet) + Omegaven 10% ($\omega 3:\omega 6$ ratio was 1:4) vs EN (standard diet)	FO 28-d Hospital 7/31 (23) 9/31 (29)	Standard 28-d Hospital 13/30 (43) 14/30 (47)	NR	NR
23) Khor 2011	ICU patients with severe sepsis/septic shock N = 27 (of 28)	C.Random: yes ITT: No Blinding: double (8)	EN and/or oral diet supplemented with 100 ml 10% Omegaven (10g refined FO, EPA 12.5-28.2 g/L, DHA 14.4-30.9 g/L) vs. 100 ml 0.9% normal saline + EN and/or oral diet	FO NR	Standard NR	NR	NR
24) Zhao 2011	ICU patients with sepsis N=116	C.Random: no ITT: no Blinding: no (5)	Omega-3 FO lipid emulsion (Omegaven), 100 ml qd for 5-7 days vs standard treatment	FO 28-d 8/56 (14)	Standard 28-d 11/60 (18)	NR	NR
25) Burkhart 2013	ICU Septic patients N=50	C.Random: unknown ITT: yes Blinding: single (assessor) (8)	2 ml.kg/d Omegaven vs no parenteral FO. Both groups received EN and/or PN without added fish oils at the discretion of the clinician.	FO 1-year# 13/25 (52)	Standard 1-year# 13/25 (52)	NR	NR
26) Hall 2014	ICU Septic patients N=60	C.Random: not sure ITT: yes Blinding: no (10)	Omegaven at 0.2 g FO /kg/d given at a rate of 0.05 g FO/kg/d vs no fish oils. In both group nutrition was assessed, by those patients requiring it, by the intensivists and dietitians who commenced oral, nasogastric (enteral), or parenteral nutrition as directed by the underlying pathology.	FO Hospital 28 day 4/30 (13.3) 4/30 (13.3)	Standard Hospital 28 day 9/30 (30) 8/30 (26.7)	FO 3/30 (10)	Standard 5/30 (16.7)

#Burkhart 2013: Discharged patients were contacted by telephone after 1 year or later to determine survivorship; median follow-up time 109 (9-408 days)

*Info from author

Table 1. Randomized studies evaluating type of lipids (PN) in critically ill patients (continued)

Study	LOS days		Ventilator days		Other
SO/MCT vs. SO					
1) Nijveldt 1998	MCT/SO NR	SO NR	MCT/SO 13.8 ± 2.9 (12)	SO 17.4 ± 3.0 (8)	NR
2) Lindgren 2001	MCT/SO ICU 26±6 (9)	SO ICU 20±5 (11)	MCT/SO NR	SO NR	MCT/SO SO Adverse effects 6/9 (67) 4/11 (36.3) Nitrogen balance at day 3 2.6 ± 5.6 gms -11.7 ± 4.8 gms; p=0.061 Nitrogen balance at day 6 1.6±11.6 gms -29.3±11.1 gms; p=0.08
3) Garnacho-Montero 2002	MCT/SO ICU 16.6 ± 6.1 (35)	SO ICU 15.8 ± 7 (37)	MCT/SO NR	SO NR	MCT/SO SO Retinol binding protein 1.7 ± 1 0.8 ± 0.6 Nitrogen balance 14.2 ± 2.9 11.6 ± 4
4) Iovinelli 2007	MCT/SO NR	SO NR	MCT/SO 10.6 ± 3.0 (7)	SO 13.4 ± 3.5 (7)	MCT/SO SO Time before weaning 52 ± 36 hrs 127 ± 73 hrs
SO/OO vs. SO/MCT or SO					
5) Garcia-de-Lorenzo 2005	OO/SO ICU 32.9 ± 10.61 ^a (11) Hospital 57 ± 15.26 ^a (11)	MCT/SO ICU 41.8 ± 16.57 ^a (11) Hospital 64.9 ± 27.20 ^a (11)	OO/SO 11.0 ± 11.94 ^a (11)	MCT/SO 13.0 ± 16.25 ^a (11)	OO/SO MCT/SO Multiple organ dysfunction score 11.0 ± 3.6 13.0 ± 4.9
6) Pontes-Arruda 2013	OO/SO ICU 12 (7-17) Hospital 21 (15-25)	MCTSO ICU 11 (5-14) Hospital 18 (13-23)	NR	NR	OO/SO MCTSO Nutritional Intake Lipids (g/day) 66 (61-73) 61 (54-67) Days on PN 12 (8-15) 11 (7-15) Dextrose (g/day) 288 (275-303) 281 (273-301) AAs (g/day) 87 (84-90) 87 (83-92)

7) Umpierrez 2012	OO/SO ICU 17 ± 18 (51) Hospital 40.8 ± 36 (51)	SO ICU 15.2 ± 14 (49) Hospital 46.7 ± 48 (49)	NR	NR	OO/SO 22 ± 6 SO 22 ± 5 Total Energy Intake (kcal/kg)
FO containing PN vs. SO or SO/MCT or SO/OO					
8) Grecu 2003*	FO/SO ICU 3.32 ± 1.48 (8) Hospital 11.68 ± 2.04 (28)	SO ICU 9.28 ± 3.08 (7) Hospital 20.46 ± 3.27 (26)	FO/SO 2.83 ± 1.62 (8)	SO 5.23 ± 2.80 (7)	FO/SO 2/28 (7) SO 8/26 (31) Patients undergoing reoperation for septic episode
9) Friesecke 2008	FO/MCT/SO ICU 28 ± 25 (83)	MCT/SO ICU 23 ± 20 (82)	FO/MCT/SO 22.8 ± 22.9 (83)	MCT/SO 20.5 ± 19.0 (82)	FO/MCT/SO 6/83 (7) MCT/SO 4/82 (5) Urinary Tract Infections Catheter-related infections 1/83 (1) 3/83 (4) Total EN Energy Intake (kcal/kg) 22.2 ± 5.5 21.6 ± 5.6
10) Guo 2008	FO ICU 21.1±2.9	SO ICU 28.4±4.2	NR	NR	NR
11) Qu 2009	NR	NR	NR	NR	NR
12) Wang 2009	NR	NR	NR	NR	FO 3/28 (11) SO 6/28 (21) Surgery of infected pancreatic necrosis
13) Barbosa 2010	FO/MCT/SO ICU 12 ± 14.4 ^a (13) Hospital 22 ± 25.2 ^a (13)	MCT/SO ICU 13 ± 12.6 ^a (10) Hospital 55 ± 50.6 ^a (10)	FO/MCT/SO 10 ± 14.4 (13)	MCT/SO 11 ± 12.64 (10)	FO/MCT/SO 2057± 418 kcals MCT/SO 1857 ± 255 kcals PO2/FiO2 ratio at day 6 331±71 vs 245±107 (p=0.047)
14) Sabater 2011	NR	NR	NR	NR	NR
15) Grau Carmona 2014	FO/MCT/SO ICU 18.9±15.5 (81)	MCT/SO ICU 21.8±20.9 (78)	FO/MCT/SO 8.4±6.6 (67)	MCT/SO 9.2±6.9 (64)	FO/MCT/SO 1.04 ± 0.12 MCT/SO 1.05 ± 0.13 Parenteral lipid intake [(g/kg BW)/d] PN kcal

	Hospital 41.1±41.0 (81)	Hospital 42.5±28.5 (78)			1,737 ± 353	1,782 ± 312
16) Gultekin 2014	FO/OO/SO Hospital 31.6 ± 17.2 (16)	OO/SO Hospital 30.6 ± 17.2 (16)	NR	NR	FO/OO/SO Kcal/kg/day 27.5±1.5	OO/SO 15.8±1.5 g protein/kg/d 1.1±0.1
17) Wang 2014	Routine TPN+ FO 7.75±1.90 (25)	Routine TPN 10.03±2.15 (28); p<0.01	Routine TPN+ FO 2.43±1.06 (25)	Routine TPN 2.94±1.37 (28); p>0.05	Routine TPN+ FO Days on CRRT 3.15±1.98	Routine TPN 3.83±1.32; p>0.05
18) Chen 2017a	NR	NR	NR	NR	NR	NR
19) Chen 2017b	FO ICU 13.8 ± 9.9	SO ICU 24.4 ± 23.2, p = 0.046	NR	NR	FO APACHE II day 7 16.1 ± 6.1	SO 21.5 ± 8, p=0.019 MARSHALL score 6.2 (2.5) 8.6 (4.3), p=0.026
20) Donoghue 2019	SMOF ICU 9.5 ± 7.09	SO ICU 10.7 ± 7.6; p= 0.49	SMOF 1.24 ± 0.83	SO 0.88 ± 1.63; p=0.39	SMOF Day 3 maximum intake of PN before start of EN Energy, kcal 2249.9 ± 386.1 vs. 2222.6 ± 352.4; p=0.76	SO Protein, g/kg 1.33±0.16 vs. 1.39 ± 0.2; p=0.16
21) Singer 2021	FO ICU 23 (15-28) Hospital 33 (24-57)	MCT/SO ICU 24 (17-36); p=0.68 Hospital 39 (26-65); p=0.45	Ventilation free days, Omega 3 FO 9.2 ± 9.4 (48)	Ventilation free days, MCT/SO 7.3 ± 9.3 (47); p=0.33	FO Days on supplemental PN and EN PN 7.1±4.0, EN 12.3±7.6 vs. PN 7.9±4.7, EN 10.6±6.7 Change in blood oxygenation, Day 1-4 -1.3±83.7 vs. 13.3± 86.1; p=0.78 Catecholamine treatment free days Higher in Omega 3 EN+PN group; p=0.05 % patients tolerating EN alone at day 6 51% vs. 29.8%; p=0.034	MCT/SO
Stand-alone FO emulsion supplement vs. standard care						
22) Gupta 2011	FO ICU 15.27 ± 9.54 (31) Hospital 19.00 ± 13.26 (31)	Standard ICU 13.70 ± 11.56 (30) Hospital 19.30 ± 16.65 (30)	FO 13.39 ± 8.30 (31)	Standard 11.30 ± 10.44 (30)		
23) Khor 2011	FO ICU 10.3 ± 7.4 (14)	Standard ICU 8.4 ± 5.7 (13)	NR	NR	NR	NR

	Hospital 19.6 ± 6.5 (14)	Hospital 17.5 ± 5.3 (13)				
24) Zhao 2011	FO ICU 8.0 ± 2.02 (56)	Standard ICU 10.97 ± 2.02 (60)	NR	NR	NR	
25) Burkhart 2013	FO ICU 5 (3-22)	Standard ICU 6 (2-33)	NR	NR	FO Subsyndromal delirium 5 (25) Sepsis associated delirium 15 (75)	Standard 6(29) 15 (71)
26) Hall 2014	FO ICU 8.8±7.7 Hospital 26.7±18.2	Standard ICU 12.3±12.4 Hospital 33.5±30.4	NR (reported as free ventilator days)	NR (reported as free ventilator days)	NR	

C.Random: concealed randomization

ITT: intent to treat

NR: not reported

* data obtained from author, 8 out of 28 in Omegaven and 7 out of 26 in LCT group were in ICU

^a converted Standard Error Mean (SEM) to Standard deviation (SD)

MCT: medium chain triglycerides

LCT: long chain triglycerides

† hospital mortality unless specified

‡ number of patients with infections unless specified

Table 2: Dose of Fish Oils in PN Trials

Study	Design	Patients	N	Nutrition / Products	Dose of FO (g/kg/d or g/d)
Greco 2003 (abstract)	RCT, DB	Abdominal sepsis	54 (15 ICU)	PN LCT + Omegaven vs. LCT	0.15 g/kg/d
Friesecke 2008	RCT, DB	115 SIRS, 52 non-SIRS	166	PN Lipofundin MCT + Omegaven vs. Lipofundin MCT	0.15 g/kg/d
Guo 2008	RCT	Septic ICU pts APACHE II>12	88	PN 20% ILE + ω -3 vs. 20% ILE	10 g/day
Wang 2009	RCT, DB	Severe acute pancreatitis in ICU	56	PN Lipovenoes + Omegaven vs. Lipovenoes	0.16 on day 1, then 0.2 g/kg/d
Barbosa 2010	RCT, SB	ICU, SIRS or sepsis requiring PN	25	PN Lipoplus vs. Nutriflex lipid special	0.09 g/kg/d
Sabater 2011	RCT	ARDS requiring MV and PN	16	PN Lipoplus 20% vs Intralipid 20%	0.12 g/kg/h
Grau Carmona 2014	RCT, DB	Medical and surgical ICU pts requiring TPN	175	PN Lipoplus vs. Lipofundin MCT	0.1 g/kg/d
Gultekin 2014	RCT, DB	Severe sepsis or septic shock	58	PN ClinOleic + Omegaven vs. ClinOleic	10 g/day
Wang 2014	RCT	Abdominal sepsis and require TPN for \geq 5 days	53	PN Routine PN + Omegaven to replace part of routine PN lipid emulsion for 5 days vs Routine PN	0.2 g/kg/day
Chen 2017a	RCT, SB	ICU pts with SIRS and ID	78	PN SO + Omegaven vs. SO	10 g/day
Chen 2017b	RCT, SB	ICU pts with SIRS and ID	48	PN Standard PN + Omegaven vs. Standard PN	10 g/day
Singer 2021	RCT, DB	ICU pts on MV	100	sPN+ EN Nutriflex Omega special + Oxepa vs. Nutriflex Lipid + Pulmocare	<0.05 g/kg/d on d1-5 >0.05 g/kg/d from d6
Donoghue 2019	RCT, DB	Critically ill, SIRS, sepsis and/ or ARDS	75	PN SMOFipid vs. Intralipid	0.09 – 0.19

Overall mortality

Figure 1.1: Overall Mortality in Trials Using an Omega-6 Reducing Strategy




Figure 1.2 Overall Mortality in Trials Using an Omega-6 Reducing strategy: Sensitivity Analyses Without Singer 2021




Figure 1.3 Overall Mortality in Trials Using an Omega-6 Reducing Strategy: Subgroup Analyses of SO/MCT or SO/OO vs. SO and FO vs SO or SO/MCT or SO/OO




Figure 1.4 Overall Mortality in PN Trials Using Fish Oils: Subgroup Analyses of Omegaven vs. Other Fish Oil Emulsions




Figure 1.5 Overall Mortality in PN Trials Using Fish Oils: Subgroup Analyses of Omegaven vs. Other Fish Oil Emulsions: Sensitivity Analyses Without Singer 2021





Figure 1.6 Overall Mortality in Trials Comparing Stand-Alone Fish Oil Emulsions to Standard of Care



28-day mortality

Figure 2.1: 28-day Mortality in Trials Using an Omega-6 Reducing Strategy




Figure 2.2 28-day Mortality in Trials Using an Omega-6 Reducing strategy: Sensitivity Analyses Without Singer 2021




Figure 2.3 28-day Mortality in Trials Using an Omega-6 Reducing Strategy: Subgroup Analyses of SO/MCT or SO/OO vs. SO and FO vs SO or SO/MCT or SO/OO




Figure 2.4 28-day Mortality in PN Trials Using Fish Oils: Subgroup Analyses of Omegaven vs. Other Fish Oil Emulsions




Figure 2.5 28-day Mortality in PN Trials Using Fish Oils: Subgroup Analyses of Omegaven vs. Other Fish Oil Emulsions: Sensitivity Analyses Without Singer 2021





Figure 2.6 28-day Mortality in Trials Comparing Stand-Alone Fish Oil Emulsions to Standard of Care



ICU Mortality

Figure 3.1 ICU Mortality in Trials Using an Omega-6 Reducing Strategy




Figure 3.2 ICU Mortality in Trials Using an Omega-6 Reducing Strategy: Subgroup Analyses of SO/MCT or SO/OO vs. SO and FO vs SO or SO/MCT or SO/OO




Figure 3.3 ICU Mortality in PN Trials Using Fish Oils: Subgroup Analyses of Omegaven vs. Other Fish Oil Emulsions




Figure 3.4 ICU Mortality in Trials Comparing Stand-Alone Fish Oil Emulsions to Standard of Care
None reported ICU mortality

Hospital Mortality

Figure 4.1 Hospital Mortality in Trials Using an Omega-6 Reducing Strategy




Figure 4.2 Hospital Mortality in Trials Using an Omega-6 Reducing Strategy: Subgroup Analyses of SO/MCT or SO/OO vs. SO and FO vs SO or SO/MCT or SO/OO




Figure 4.3 Hospital Mortality in PN Trials Using Fish Oils: Subgroup Analyses of Omegaven vs. Other Fish Oil Emulsions




Figure 4.4 Hospital Mortality in Trials Comparing Stand-Alone Fish Oil Emulsions to Standard of Care




Figure 5.1 Overall Infections in Trials Using an Omega-6 Reducing strategy




Figure 5.2 Overall Infections in PN Trials Using Fish Oils: Subgroup Analyses of Omegaven vs. Other Fish Oil Emulsions




Figure 6.1 Hospital LOS in Trials Using an Omega-6 Reducing Strategy




Figure 6.2 Hospital LOS in PN Trials Using Fish Oils: Subgroup Analyses of Omegaven vs. Other Fish Oil Emulsions




Figure 6.3 Hospital LOS in Trials Comparing Stand-Alone Fish Oil Emulsions to Standard of Care




Figure 7.1 ICU LOS in Trials Using an Omega-6 Reducing Strategy




Figure 7.2 ICU LOS in Trials Using an Omega-6 Reducing Strategy: Subgroup Analyses of SO/MCT or SO/OO vs. SO and FO vs SO or SO/MCT or SO/OO




Figure 7.3. ICU LOS in PN Trials Using Fish Oils: Subgroup Analyses of Omegaven vs. Other Fish Oil Emulsions




Figure 7.4. ICU LOS in Trials Comparing Stand-alone Fish Oil Emulsions to Standard of Care




Figure 8.1 Ventilator Days in Trials Using an Omega-6 Reducing Strategy




Figure 8.2 Ventilator Days in Trials Using an Omega-6 Reducing Strategy: Subgroup Analyses of SO/MCT or SO/OO vs. SO and FO vs SO or SO/MCT or SO/OO




Figure 8.3 Ventilator Days in PN Trials Using Fish Oils: Subgroup Analyses of Omegaven vs. Other Fish Oil Emulsions





Figure 8.4 Ventilator Days in Trials Comparing Stand-Alone Fish Oil Emulsions to Standard of Care



References

Included Studies

Soybean Oil

1. Nijveldt RJ, Tan AM, Prins HA, de Jong D, van Rij GL, Wesdorp RI, van Leeuwen PA. Use of a mixture of medium-chain triglycerides and longchain triglycerides versus long-chain triglycerides in critically ill surgical patients: a randomized prospective double-blind study. *Clin Nutr.* 1998 Feb;17(1):23-9.
2. Lindgren BF, Ruokonen E, Magnusson-Borg K, Takala J. Nitrogen sparing effect of structured triglycerides containing both medium-and long-chain fatty acids in critically ill patients; a double blind randomized controlled trial. *Clin Nutr.* 2001 Feb;20(1):43-8.
3. Garnacho-Montero J, Ortiz-Leyba C, Jiménez-Jiménez FJ, Garcia-Garmendia JL, Jiménez-Jiménez LM, Garnacho-Montero MC, Barrero-Almodóvar A.. Clinical and metabolic effects of two lipid emulsions on the parenteral nutrition of septic patients. *Nutrition.* 2002 Feb;18(2):134-8
4. Lovinelli G, Marinangeli F, Ciccone A, Ciccozzi A, Leonardis M, Paladini A, Varrassi G. Parenteral nutrition in ventilated patients with chronic obstructive pulmonary disease: long chain vs medium chain triglycerides. *Minerva Anestesiol.* 2007 Jan-Feb;73(1-2):65-76.

Olive Oil containing emulsions

5. García-de-Lorenzo A, Denia R, Atlan P, Martínez-Ratero S, Le Brun A, Evard D, Bereziat G. Parenteral nutrition providing a restricted amount of linoleic acid in severely burned patients: a randomised double-blind study of an olive oil-based lipid emulsion v. medium/long-chain triacylglycerols. *Br J Nutr.* 2005 Aug;94(2):221-30.
6. "Pontes-Arruda A, Dos Santos MC, Martins LF, González ER, Kliger RG, Maia M, Magnan GB; EPICOS Study Group. Influence of parenteral nutrition delivery system on the development of bloodstream infections in critically ill patients: an international, multicenter, prospective, open-label, controlled study--EPICOS study. *JPEN J Parenter Enteral Nutr.* 2012 Sep;36(5):574-86. Epub 2012 Jan 23. PubMed PMID: 22269899."
7. Umpierrez GE, Spiegelman R, Zhao V, Smiley DD, Pinzon I, Griffith DP, et al. A double-blind, randomized clinical trial comparing soybean oil-based versus olive oil-based lipid emulsions in adult medical-surgical intensive care unit patients requiring parenteral nutrition. *Crit Care Med* 2012;40(6):1792-8.

Fish Oil containing PN

8. Grecu I, Mirea L, Grintescu I. Parenteral fish oil supplementation in patients with abdominal sepsis. *Clinical Nutrition* 2003 [abstract]
9. Friesecke S, Lotze C, Köhler J, Heinrich A, Felix SB, Abel P. Fish oil supplementation in the parenteral nutrition of critically ill medical patients: a randomised controlled trial. *Intensive Care Med.* 2008 Aug;34(8):1411-20.
10. Guo Y. ω -3 duo bu baohe zhifangsuān dui nong du zheng huanzhe linchuang liaoxiao ji yuhou de yingxiang (Effect of omega-3 polyunsaturated fatty acids on clinical efficacy and prognosis in patients with sepsis). *Shanxi Med J.* 2008;37(15):727-9

11. Qu A, Xu L. Yuyou zhifang ru dui nong du zheng huanzhe mianyi gongneng de tiaojie zuoyong guan cha (Regulation of fish oil fat emulsion on immune function in patients with sepsis). *Shandong Med J*. 2009;49(6):13–5.
12. Wang X, Li W, Zhang F, Pan L, Li N, Li J. Fish oil-supplemented parenteral nutrition in severe acute pancreatitis patients and effects on immune function and infectious risk: a randomized controlled trial. *Inflammation*. 2009 Oct;32(5):304-9. PubMed PMID: 19568921
13. Barbosa VM, Miles EA, Calhau C, Lafuente E, Calder PC. Effects of a fish oil containing lipid emulsion on plasma phospholipid fatty acids, inflammatory markers, and clinical outcomes in septic patients: a randomized, controlled clinical trial. *Crit Care*. 2010;14(1):R5. PubMed PMID: 20085628; PubMed Central PMCID: PMC2875515
14. Sabater J, Masclans JR, Sacanell J, Chacon P, Sabin P, Planas M. Effects of an omega-3 fatty acid-enriched lipid emulsion on eicosanoid synthesis in acute respiratory distress syndrome (ARDS): a prospective, randomized, double-blind, parallel group study. *Nutr Metab (Lond)*. 2011;8(1):22.
15. Grau-Carmona T, Bonet-Saris A, García-de-Lorenzo A, Sánchez-Alvarez C, Rodríguez-Pozo A, Acosta-Escribano J, Miñambres E, Herrero-Meseguer JI, Mesejo A. Influence of n-3 Polyunsaturated Fatty Acids Enriched Lipid Emulsions on Nosocomial Infections and Clinical Outcomes in Critically Ill Patients: ICU Lipids Study. *Crit Care Med*. 2015 Jan;43(1):31-9.
16. Gultekin G, Sahin H, Inanc N, Uyanik F, Ok E. Impact of Omega-3 and Omega-9 fatty acids enriched total parenteral nutrition on blood chemistry and inflammatory markers in septic patients. *Pak J Med Sci*. 2014 Mar;30(2):299-304.
17. Wang X, Tong F, Chen H, Huo SH, Liu F, Liu ZK, et al. Clinical effect of omega-3 polyunsaturated fatty acid on patients with abdominal sepsis. *Clinical Focus*. 2014;29:502–4,8.
18. **Chen 2017 A** - Chen H, Wang W, Hong C, Zhang M, Hong Y, Wang S, Zhang H. Omega-3 Fish Oil Reduces Mortality Due to Severe Sepsis with Acute Gastrointestinal Injury Grade III. *Pharmacogn Mag*. 2017 Jul-Sep;13(51):407-412. 2017 Jul 19.
19. **Chen 2017 B**- Chen H, Wang W, Hong Y, Zhang H, Hong C, Liu X. Single-blinded, randomized, and controlled clinical trial evaluating the effects of Omega-3 fatty acids among septic patients with intestinal dysfunction: A pilot study. *Exp Ther Med*. 2017;14(2):1505-1511. doi:10.3892/etm.2017.4680
20. Donoghue, V.; Schleicher, G. ; Spruyt, M. G. .; Malan, L.; Nel, D. .; Calder, P. .; Blaauw, R. Four-Oil Intravenous Lipid Emulsion Effect on Plasma Fatty Acid Composition, Inflammatory Markers and Clinical Outcomes in Acutely Ill Patients: A Randomised Control Trial (Foil Fact). *Clinical nutrition (Edinburgh, Scotland)* 2019, 38 (6), 2583–2591. <https://doi.org/10.1016/j.clnu.2018.12.010>.
21. Singer, P. Bendavid, I., Mesilati-Stahy, R. et. al. Enteral and supplemental parenteral nutrition enriched with omega-3 polyunsaturated fatty acids in intensive care patients: A randomized, controlled, double-blind clinical trial. *Clinical Nutrition* 2021 (in press).

Stand alone Fish Oil

22. Gupta A, Govil D, Bhatnagar S, et. al. Efficacy and safety of parenteral omega 3 fatty acids in ventilated patients with acute lung injury. *InJCCM* April-June 2011, Vol 5, 108-113.
23. Khor BS, Liaw SJ, Shih HC, Wang LS. Randomized, double blind, placebo-controlled trial of fish-oil-based lipid emulsion infusion for treatment of critically ill patients with severe sepsis. *Asian J Surg*. 2011;34(1):1-10. doi:10.1016/S1015-9584(11)60011-0
24. Zhao K, Zhou W, Bo C. ω-3 yuyou zhifang ru dui nong du zheng huanzhe de linchuang liaoxiao (Clinical efficacy of omega-3 fish oil fat emulsion in patients with sepsis). *Shandong Med J*. 2011;51(16):102.

25. Burkhart CS, Dell-Kuster S, Siegemund M, Pargger H, Marsch S, Strebel SP, Steiner LA. Effect of n-3 fatty acids on markers of brain injury and incidence of sepsis-associated delirium in septic patients. *Acta Anaesthesiol Scand*. 2014 Mar 24. [Epub]
26. Hall TC, Bilku DK, Al-Leswas D, Neal CP, Horst C, Cooke J, Metcalfe MS, Dennison AR. A randomized controlled trial investigating the effects of parenteral fish oil on survival outcomes in critically ill patients with sepsis: a pilot study. *JPEN J Parenter Enteral Nutr*. 2015 Mar;39(3):301-12. doi: 10.1177/0148607113518945. Epub 2014 Jan 9. PMID: 24408035.

Exclusion list

#	Reason excluded	Citation
1	Duplicate of Wang 2009	Wang X, Li W, Li N, Li J. Omega-3 fatty acids-supplemented parenteral nutrition decreases hyperinflammatory response and attenuates systemic disease sequelae in severe acute pancreatitis: a randomized and controlled study. JPEN J Parenter Enteral Nutr. 2008;32(3):236-241. doi:10.1177/0148607108316189
2	Elective surgery patients	Martínez-Lozano Aranaga F, Gómez Ramos MJ, Sánchez Álvarez MDC. Eficacia y seguridad de dos emulsiones lipídicas de nutrición parenteral en pacientes críticos posquirúrgicos: Clinoleic® frente a SMOFlipid® [Effectiveness and safety of two lipid emulsions for parenteral nutrition in postsurgical critically ill patients: Clinoleic® versus SMOFlipid®]. Nutr Hosp. 2021;38(1):5-10. doi:10.20960/nh.03077
3	Elective surgery patients	Wachtler P, König W, Senkal M, Kemen M, Köller M. Influence of a total parenteral nutrition enriched with omega-3 fatty acids on leukotriene synthesis of peripheral leukocytes and systemic cytokine levels in patients with major surgery. J Trauma. 1997 Feb;42(2):191-8.
4	Elective surgery patients	Chambrier C, Guiraud M, Gibault JP, Labrosse H, Boulétreau P. Medium- and long-chain triacylglycerols in postoperative patients: structured lipids versus a physical mixture. Nutrition. 1999 Apr;15(4):274-7.
5	Elective surgery patients	Hailer S, Jauch KW, Wolfram G. Influence of different fat emulsions with 10 or 20% MCT/LCT or LCT on lipoproteins in plasma of patients after abdominal surgery. Ann Nutr Metab. 1998;42(3):170-80.
6	Elective surgery patients	Linseisen J, Hoffmann J, Lienhard S, Jauch KW, Wolfram G. Antioxidant status of surgical patients receiving TPN with an omega-3-fatty acid-containing lipid emulsion supplemented with alpha-tocopherol. Clin Nutr. 2000 Jun;19(3):177-84.
7	Elective surgery patients	Heller AR, Fischer S, Rossel T, Geiger S, Siegert G, Ragaller M, et al. Impact of n-3 fatty acid supplemented parenteral nutrition on haemostasis patterns after major abdominal surgery. The British journal of nutrition. 2002;87 Suppl 1:S95-101.
8	Elective surgery patients	Kuse ER, Kotzerke J, Müller S, Nashan B, Lück R, Jaeger K. Hepatic reticuloendothelial function during parenteral nutrition including an MCT/LCT or LCT emulsion after liver transplantation - a double-blind study. Transpl Int. 2002 Jun;15(6):272-7. Epub 2002 Apr 30.
9	Elective surgery patients	Weiss G, Meyer F, Matthies B, Pross M, Koenig W, Lippert H. Immunomodulation by perioperative administration of n-3 fatty acids. The British journal of nutrition 2002;87 Suppl 1:S89-94.
10	Elective surgery patients	Grau T, Ruiz de Adana JC, Zubillaga S, Fuerte S, Girón C. [Randomized study of two different fat emulsions in total parenteral nutrition of malnourished surgical patients;effect of infectious morbidity and mortality] [Article in Spanish]. Nutr Hosp. 2003 May-Jun;18(3):159-66.
11	Elective surgery patients	Heller AR, Rössel T, Gottschlich B, Tiebel O, Menschikowski M, Litz RJ, Zimmermann T, Koch T. Omega-3 fatty acids improve liver and pancreas function in postoperative cancer patients. Int J Cancer. 2004 Sep 10;111(4):611-6.
12	Elective surgery patients	Chen FM, Wang JY, Sun LC, Juang RF, Huang TJ, Hsieh JS. Efficacy of medium-chain triglycerides compared with long-chain triglycerides in total parenteral nutrition in patients with digestive tract cancer undergoing surgery. Kaohsiung J Med Sci. 2005 Nov;21(11):487-94.
13	Elective surgery patients	Senkal M, Geier B, Hannemann M, Deska T, Linseisen J, Wolfram G, Adolph M. Supplementation of omega-3 fatty acids in parenteral nutrition beneficially alters phospholipid fatty acid pattern. JPEN J Parenter Enteral Nutr. 2007 Jan-Feb;31(1):12-7.

14	Elective surgery patients	Wendel M et al. Impact of total parenteral nutrition including omega-3 fatty acids on the regulation of plasma lipoproteins and glycemic control after major abdominal surgery. <i>E-SPEN</i> . 2007. 2:e103-e110
15	Elective surgery patients	Wichmann MW, Thul P, Czarnetzki HD, Morlion BJ, Kemen M, Jauch KW. Evaluation of clinical safety and beneficial effects of a fish oil containing lipid emulsion (Lipoplus, MLF541): data from a prospective, randomized, multicenter trial. <i>Crit Care Med</i> . 2007;35(3):700-706. doi:10.1097/01.CCM.0000257465.60287.AC
16	Elective surgery patients	Berger MM, Tappy L, Revelly JP, Koletzko BV, Gepert J, Corpataux JM, Cayeux MC, Chiolo RL. Fish oil after abdominal aorta aneurysm surgery. <i>Eur J Clin Nutr</i> . 2008 Sep;62(9):1116-22. Epub 2007 May 30. Erratum in: <i>Eur J Clin Nutr</i> . 2009 Feb;63(2):302.
17	Elective surgery patients	Shan Y-S et al. Evaluation of the stability and safety of Venolipid MCT/LCT 20% administered by an all-in-one system in patients after major gastrointestinal surgery. <i>E-SPEN</i> . 2008(3):e135-e141
18	Elective surgery patients	Puiggròs C, Sánchez J, Chacón P, Sabin P, Roselló J, Bou R, Planas M. Evolution of lipid profile, liver function, and pattern of plasma fatty acids according to the type of lipid emulsion administered in parenteral nutrition in the early postoperative period after digestive surgery. <i>JPEN J Parenter Enteral Nutr</i> . 2009 Sep-Oct;33(5):501-12.
19	Elective surgery patients	Badía-Tahull MB, Llop-Talaverón JM, Leiva-Badosa E, Biondo S, Farran-Teixidó L, Ramón-Torrell JM, Jódar-Masanes R. A randomised study on the clinical progress of high-risk elective major gastrointestinal surgery patients treated with olive oil-based parenteral nutrition with or without a fish oil supplement. <i>Br J Nutr</i> . 2010;104(5):737-41.
20	Elective surgery patients	Makay O, Kaya T, Firat O, Sozbilen M, Caliskan C, Gezer G, et al. omega-3 Fatty acids have no impact on serum lactate levels after major gastric cancer surgery. <i>JPEN Journal of parenteral and enteral nutrition</i> 2011;35(4):488-92.
21	Elective surgery patients	Zhu MW, Tang DN, Hou J, Wei JM, Hua B, Sun JH, et al. Impact of fish oil enriched total parenteral nutrition on elderly patients after colorectal cancer surgery. <i>Chinese medical journal</i> . 2012;125(2):178-81.
22	Elective surgery patients	Berger MM, Delodder F, Liaudet L, Tozzi P, Schlaepfer J, Chiolo RL, Tappy L. Three short perioperative infusions of n-3 PUFAs reduce systemic inflammation induced by cardiopulmonary bypass surgery: a randomized controlled trial. <i>Am J Clin Nutr</i> . 2013 Feb;97(2):246-54. doi: 10.3945/ajcn.112.046573. Epub 2012 Dec 26. PubMed PMID: 23269816.
23	Elective surgery patients	Metry AA, Abdelaal W, Ragaei M, Refaat M, Nakhla G. SMOFlipid versus Intralipid in Postoperative ICU Patients. <i>Enliven Archive</i> . 2014;1(6):1-8.
24	Elective surgery patients	Demirer S, Sapmaz A, Karaca AS, et al. Effects of postoperative par- enteral nutrition with different lipid emulsions in patients undergoing major abdominal surgery. <i>Ann Surg Treat Res</i> . 2016;91(6):309-315.
25	Elective surgery patients	Gong Y, Liu Z, Liao Y, et al. Effectiveness of ω -3 Polyunsaturated Fatty Acids Based Lipid Emulsions for Treatment of Patients after Hepatectomy: A Prospective Clinical Trial. <i>Nutrients</i> . 2016;8(6):357. Published 2016 Jun 17. doi:10.3390/nu8060357
26	Elective surgery patients	Fiaccadori E, Tortorella G, Gonzi GL, Beghi C, Albertini D, Pincolini S, Belli L, Avogaro, A. Hemodynamic and respiratory effects of medium-chain and long-chain triglyceride fat emulsions: A prospective, randomized study. <i>Rivista Italiana di Nutrizione Parenterale ed Enterale</i> 1997;15(1):6-14
27	EN only	Yan H, Huang XH, Xiao KJ, et al. <i>Zhonghua Shao Shang Za Zhi</i> . 2003;19(4):202-205.

28	EN only	Bernier J, Jobin N, Emptoz-Bonneton A, Pugeat MM, Garrel DR. Decreased corticosteroid-binding globulin in burn patients: relationship with interleukin-6 and fat in nutritional support. <i>Crit Care Med.</i> 1998;26(3):452-460. doi:10.1097/00003246-199803000-00014
29	EN only	Adams S, Yeh YY, Jensen GL. Changes in plasma and erythrocyte fatty acids in patients fed enteral formulas containing different fats. <i>JPEN J Parenter Enteral Nutr.</i> 1993 Jan-Feb;17(1):30-4.
30	EN only	Gogos CA, Ginopoulos P, Salsa B, Apostolidou E, Zoumbos NC, Kalfarentzos F. Dietary omega-3 polyunsaturated fatty acids plus vitamin E restore immunodeficiency and prolong survival for severely ill patients with generalized malignancy: a randomized control trial. <i>Cancer.</i> 1998 Jan 15;82(2):395-402.
31	EN only	Najmi M, Vahdat Shariatpanahi Z, Tolouei M, Amiri Z. Effect of oral olive oil on healing of 10-20% total body surface area burn wounds in hospitalized patients. <i>Burns.</i> 2015 May;41(3):493-6.
32	EN only	Theilla, M.; Singer, P.; Cohen, J.; Dekeyser, F. A diet enriched in eicosapentanoic acid, gamma-linolenic acid and antioxidants in the prevention of new pressure ulcer formation in critically ill patients with acute lung injury: A randomized, prospective, controlled study.
33	EN only	Hosny M, Nahas R, Ali S, Elshafei SA, Khaled H. Impact of oral omega-3 fatty acids supplementation in early sepsis on clinical outcome and immunomodulation. <i>Egypt J Crit Care Med.</i> 2013;1:119–26.
34	Lipid emulsion only infused for 12 hrs on one day, followed by standard PN	Sabater J, Masclans JR, Sacanell J, Chacon P, Sabin P, Planas M. Effects on hemodynamics and gas exchange of omega-3 fatty acid-enriched lipid emulsion in acute respiratory distress syndrome (ARDS): a prospective, randomized, double-blind, parallel group study. <i>Lipids in health and disease.</i> 2008;7:39.
35	Lipid vs no lipid	Taftachi F, Sanaei-Zadeh H, Sepehrian B, Zamani N. Lipid emulsion improves Glasgow coma scale and decreases blood glucose level in the setting of acute non-local anesthetic drug poisoning--a randomized controlled trial. <i>European review for medical and pharmacological sciences</i> 2012;16 Suppl 1:38-42.
36	Lipid vs no lipid	Battistella FD, Widergren JT, Anderson JT, Siepler JK, Weber JC, MacColl K. A prospective, randomized trial of intravenous fat emulsion administration in trauma victims requiring total parenteral nutrition. <i>J Trauma.</i> 1997;43(1):52-60. doi:10.1097/00005373-199707000-00013
37	Lipid vs no lipid	Lekka ME, Liokatis S, Nathanail C, Galani V, Nakos G. The impact of intravenous fat emulsion administration in acute lung injury. <i>Am J Respir Crit Care Med.</i> 2004;169(5):638-644. doi:10.1164/rccm.200305-6200C
38	Lipid vs no lipid	Adolph M, Hailer S, Echart J. Serum phospholipid fatty acids in severely injured patients on total parenteral nutrition with medium chain/long chain triglyceride emulsions. <i>Ann Nutr Metab.</i> 1995;39(4):251-260. doi:10.1159/000177870
39	No clinical outcomes	Calon B, Pottecher T, Frey A, Ravanello J, Otteni JC, Bach AC. Long-chain versus medium and long-chain triglyceride-based fat emulsion in parental nutrition of severe head trauma patients. <i>Infusionstherapie.</i> 1990 Oct;17(5):246-8.
40	No clinical outcomes	Hwang TL, Huang SL, Chen MF. Effects of intravenous fat emulsion on respiratory failure. <i>Chest</i> 1990;97(4):934-8.
41	No clinical outcomes	Jarnberg P-O. Liposyn versus intralipid: A comparative study of two lipid emulsion in critically ill patients receiving total parenteral nutrition. 1991 50(1):38-44
42	No clinical outcomes	Diboune M, Ferard G, Ingenbleek Y, Tulasne PA, Calon B, Hasselmann M, Sauder P, Spielmann D, Metais P. Composition of phospholipid fatty acids in red blood cell membranes of patients in intensive care units: effects of different intakes of soybean oil, medium-chain triglycerides, and black-currant seed oil. <i>JPEN J Parenter Enteral Nutr.</i> 1992 Mar-Apr;16(2):136-41.

43	No clinical outcomes	Ball MJ. Parenteral nutrition in the critically ill: use of a medium chain triglyceride emulsion. <i>Intensive Care Med.</i> 1993;19(2):89-95.
44	No clinical outcomes	Chassard D, Guiraud M, Gauthier J, Gelas P, Berrada KR, Bouletreau P. Effects of intravenous medium-chain triglycerides on pulmonary gas exchanges in mechanically ventilated patients. <i>Crit Care Med.</i> 1994 Feb;22(2):248-51.
45	No clinical outcomes	Jeevanandam M, Holaday NJ, Voss T, Buier R, Petersen SR. Efficacy of a mixture of medium-chain triglyceride (75%) and long-chain triglyceride (25%) fat emulsions in the nutritional management of multiple-trauma patients. <i>Nutrition.</i> 1995 May-Jun;11(3):275-84.
46	No clinical outcomes	Kalfarentzos F, Kokkinis K, Leukaditi K, Maroulis J, Onoufriou A, Alexopoulos K. Comparison between two fat emulsions: Intralipid 30 cent vs intralipid 10 cent in critically ill patients. <i>Clin Nutr.</i> 1998 Feb;17(1):31-4.
47	No clinical outcomes	Masclans JR, Iglesia R, Bermejo B, Picó M, Rodriguez-Roisin R, Planas M. Gas exchange and pulmonary haemodynamic responses to fat emulsions in acute respiratory distress syndrome. <i>Intensive Care Med.</i> 1998 Sep;24(9):918-23.
48	No clinical outcomes	Smirmiotis V, Kostopanagiotou G, Vassiliou J, Arkadopoulos N, Vassiliou P, Datsis A, et al. Long chain versus medium chain lipids in patients with ARDS: effects on pulmonary haemodynamics and gas exchange. <i>Intensive care medicine</i> 1998;24(10):1029-33.
49	No clinical outcomes	Planas M, Porta I, Sagristá ML, Mora M, Padró JB, Picó M. Fatty acid composition of platelet membrane lipids after administration of two different fat emulsions in critically ill patients. <i>Intensive Care Med.</i> 1999 Apr;25(4):395-8.
50	No clinical outcomes	Manuel-y-Keenoy B, Nonneman L, De Bosscher H, Vertommen J, Schrans S, Klütsch K, De Leeuw I. Effects of intravenous supplementation with alpha-tocopherol in patients receiving total parenteral nutrition containing medium- and long-chain triglycerides. <i>Eur J Clin Nutr.</i> 2002 Feb;56(2):121-8.
51	No clinical outcomes	Schauder P, Rohn U, Schafer G, Korff G, Schenk HD. Impact of fish oil enriched total parenteral nutrition on DNA synthesis, cytokine release and receptor expression by lymphocytes in the postoperative period. <i>The British journal of nutrition.</i> 2002;87 Suppl 1:S103-10.
52	No clinical outcomes	García-de-Lorenzo A, López-Martínez J, Planas M, Chacón P, Montejo JC, Bonet A, Ortiz-Leyba C, Sánchez-Segura JM, Ordóñez J, Acosta J, Grau T, Jiménez FJ. Safety and metabolic tolerance of a concentrated long-chain triglyceride lipid emulsion in critically ill septic and trauma patients. <i>JPEN J Parenter Enteral Nutr.</i> 2003 May-Jun;27(3):208-15.
53	No clinical outcomes	Koller M, Senkal M, Kemen M, König W, Zumtobel V, Muhr G. Impact of omega-3 fatty acid enriched TPN on leukotriene synthesis by leukocytes after major surgery. <i>Clin Nutr.</i> 2003;22(1):59-64.
54	No clinical outcomes	Mayer K, Fegbeutel C, Hattar K, Sibelius U, Krämer HJ, Heuer KU, Temmesfeld-Wollbrück B, Gokorsch S, Grimminger F, Seeger W. Omega-3 vs. omega-6 lipid emulsions exert differential influence on neutrophils in septic shock patients: impact on plasma fatty acids and lipid mediator generation. <i>Intensive Care Med.</i> 2003 Sep;29(9):1472-81.
55	No clinical outcomes	Mayer K, Gokorsch S, Fegbeutel C, Hattar K, Rosseau S, Walmrath D, Seeger W, Grimminger F. Parenteral nutrition with fish oil modulates cytokine response in patients with sepsis. <i>Am J Respir Crit Care Med.</i> 2003 May 15;167(10):1321-8. Epub 2003 Feb 25.
56	No clinical outcomes	Mayer K, Meyer S, Reinholz-Muhly M, Maus U, Merfels M, Lohmeyer J, et al. Short-time infusion of fish oil-based lipid emulsions, approved for parenteral nutrition, reduces monocyte proinflammatory cytokine generation and adhesive interaction with endothelium in humans. <i>J Immunol.</i> 2003;171(9):4837-43.

57	No clinical outcomes	Antébi H, Mansoor O, Ferrier C, Tétégan M, Morvan C, Rangaraj J, Alcindor LG. Liver function and plasma antioxidant status in intensive care unit patients requiring total parenteral nutrition: comparison of 2 fat emulsions. <i>JPEN J Parenter Enteral Nutr.</i> 2004 May-Jun;28(3):142-8.
58	No clinical outcomes	Tappy L, Berger MM, Schwarz JM, Schneiter P, Kim S, Revely JP, Chioloro R. Metabolic effects of parenteral nutrition enriched with n-3 polyunsaturated fatty acids in critically ill patients. <i>Clin Nutr.</i> 2006 Aug;25(4):588-95.
59	No clinical outcomes	Piper SN, Schade I, Beschmann RB, Maleck WH, Boldt J, Röhm KD. Hepatocellular integrity after parenteral nutrition: comparison of a fish-oil-containing lipid emulsion with an olive-soybean oil-based lipid emulsion. <i>Eur J Anaesthesiol.</i> 2009 Dec;26(12):1076-82.
60	No clinical outcomes	Xiong J, Zhu S, Zhou Y, Wu H, Wang C. Regulation of omega-3 fish oil emulsion on the SIRS during the initial stage of severe acute pancreatitis. <i>J Huazhong Univ Sci Technolog Med Sci.</i> 2009 Feb;29(1):35-8.
62	No clinical outcomes	Sungurtekin H, Değirmenci S, Sungurtekin U, Oguz BE, Sabir N, Kaptanoglu B. Comparison of the effects of different intravenous fat emulsions in patients with systemic inflammatory response syndrome and sepsis. <i>Nutr Clin Pract.</i> 2011 Dec;26(6):665-71.
63	No clinical outcomes	Chen H, Cai L, Zhang X. Comparison of the effects of structured triglyceride and medium/long chain triglycerides on liver function and serum lipid in critical patients. <i>China Pharm</i> 2013;24(24).
64	No clinical outcomes	Planas M, Masclans JR, Iglesia R, et al: Eicosanoids and fat emulsions in acute respiratory distress syndrome patients. <i>Nutrition</i> 1997; 13: 202–205
65	No clinical outcomes	Lu M, Ye H, Yu S, Feng Y, Zheng F. Clinical application of the structured triglyceride and physical mixed MCT/LCT in critically ill patients. <i>Mod Med J China</i> 2012;14(7):8e10.
66	No clinical outcomes	Niu, G., Zhao, R., Gao, F., Zheng, D., Liu, X., & Wu, H. (2014). Effect of ω -3 polyunsaturated fatty acids on intestinal mucosal barrier of patients with severe acute pancreatitis.
67	No clinical outcomes	Su MS, He L, Liu ZW, Ma HX, Zhao QH, Zhang WZ. [Applied studies of structured triglycerides for parenteral nutrition in severe hemorrhagic shock patients after resuscitation] <i>Zhonghua Yi Xue Za Zhi.</i> 2012;92(12):827-830.
68	No clinical outcomes	Chacón Castro MP, Jiménez Sesé G, Salvadó Salvát J, Sabin Urquía P, Pascual Mostaza C, Planas Vilà M. Influencia de dos emulsiones grasas con distinta composición de triglicéridos sobre el metabolismo lipídico del paciente séptico [The effect of fatty emulsions with distinct triglyceride compositions on the lipid metabolism of the septic patient]. <i>Nutr Hosp.</i> 2000;15(1):13-17.
69	No clinical outcomes	Pastó L, Pellicer E, Llopart R, Gómez JA, Cardona D. Estudio comparativo de dos emulsiones lipídicas en enfermos que reciben nutrición parenteral total [Comparative study of two lipid emulsions in patients receiving total parenteral nutrition]. <i>Nutr Hosp.</i> 1991;6(1):29-33.
70	No clinical outcomes	Maeshima Y, Fukatsu K, Moriya T, et al. Influence of adding fish oil to parenteral nutrition on gut-associated lymphoid tissue. <i>JPEN J Parenter Enteral Nutr.</i> 2007;31(5):416-422. doi:10.1177/0148607107031005416
71	No clinical outcomes	Widhalm K, Kohl S, Hammerle A. The clinical application of two newly developed lipid emulsions (Solipid 20% S&E) in critically ill patients. <i>Infusionsther Transfusionsmed.</i> 1996;23(1):8-12. doi:10.1159/000223248
72	No true control group	Chen J, Yan J, Cai GL, Xu QH, Gong SJ, Dai HW, Yu YH, Li L. Structured lipid emulsion as nutritional therapy for the elderly patients with severe sepsis. <i>Chin Med J (Engl).</i> 2013 Jun;126(12):2329-32.

73	Not a omega-6 reducing strategy. Compares one LCT to another	Kari A, Hersio K, Takala J, Penttila I. Comparison of two long-chain triglyceride fat emulsions in parenteral nutrition of critically ill patients. <i>Current Therapeutic Research</i> . 1989 June;45(6):1077-87.
74	Not critically ill patients	Hutchison ML, Clemans G. Prospective trial of liposyn 20% in patients undergoing bone marrow transplantation. <i>Clinical Nutrition</i> . 1984 May;3:5-9
75	Not critically ill patients	Furukawa K, Tashiro T, Yamamori H, Takagi K, Morishima Y, Sugiura T, Otsubo Y, Hayashi N, Itabashi T, Sano W, Toyoda Y, Nitta H, Nakajima N. Effects of soybean oil emulsion and eicosapentaenoic acid on stress response and immune function after a severely stressful operation. <i>Ann Surg</i> . 1999 Feb;229(2):255-61.
76	Not critically ill patients	Klek S, Kulig J, Szczepanik AM, Jedrys J, Kolodziejczyk P. The clinical value of parenteral immunonutrition in surgical patients. <i>Acta Chir Belg</i> . 2005 Apr;105(2):175-9.
77	Not critically ill patients	Cano NJ, Saingra Y, Dupuy AM, Lorec-Penet AM, Portugal H, Lairon D, Cristol JP, Come A, Le Brun A, Atlan P, Leverve XM. Intradialytic parenteral nutrition: comparison of olive oil versus soybean oil-based lipid emulsions. <i>Br J Nutr</i> . 2006 Jan;95(1):152-9.
78	Not critically ill patients	Grimm H, Mertes N, Goeters C, Schlotzer E, Mayer K, Grimminger F, Fürst P. Improved fatty acid and leukotriene pattern with a novel lipid emulsion in surgical patients. <i>Eur J Nutr</i> . 2006 Feb;45(1):55-60.
79	Not critically ill patients	Mertes N, Grimm H, Fürst P, Stehle P. Safety and efficacy of a new parenteral lipid emulsion (SMOFlipid) in surgical patients: a randomized, double-blind, multicenter study. <i>Ann Nutr Metab</i> . 2006;50(3):253-9.
80	Not critically ill patients	Liang B, Wang S, Ye YJ, Yang XD, Wang YL, Qu J, Xie QW, Yin MJ. Impact of postoperative omega-3 fatty acid-supplemented parenteral nutrition on clinical outcomes and immunomodulations in colorectal cancer patients. <i>World J Gastroenterol</i> . 2008 Apr 21;14(15):2434-9.
81	Not critically ill patients	Wang X, Li W, Li N, Li J. Omega-3 fatty acids-supplemented parenteral nutrition decreases hyperinflammatory response and attenuates systemic disease sequelae in severe acute pancreatitis: a randomized and controlled study. <i>JPEN J Parenter Enteral Nutr</i> . 2008 May-Jun;32(3):236-41.
82	Not critically ill patients	Heidt MC, Vician M, Stracke SK, Stadlbauer T, Grebe MT, Boening A, Vogt PR, Erdogan A. Beneficial effects of intravenously administered N-3 fatty acids for the prevention of atrial fibrillation after coronary artery bypass surgery: a prospective randomized study. <i>Thorac Cardiovasc Surg</i> . 2009 Aug;57(5):276-80.
83	Not critically ill patients	Piper SN, Schade I, Beschmann R, Maleck W, Boldt J, Rohm KD. Hepatocellular integrity after parenteral nutrition: comparison of a fish-oil-containing lipid emulsion with an olive-soybean oil-based lipid emulsion. <i>European Journal of Anaesthesiology</i> . 2009;26(12):1076-82.
84	Not critically ill patients	Jiang ZM, Wilmore DW, Wang XR, Wei JM, Zhang ZT, Gu ZY, Wang S, Han SM, Jiang H, Yu K. Randomized clinical trial of intravenous soybean oil alone versus soybean oil plus fish oil emulsion after gastrointestinal cancer surgery. <i>Br J Surg</i> . 2010 Jun;97(6):804-9.
85	Not critically ill patients	Han YY, Lai SL, Ko WJ, Chou CH, Lai HS. Effects of fish oil on inflammatory modulation in surgical intensive care unit patients. <i>Nutr Clin Pract</i> . 2012 Feb;27(1):91-8. Epub 2012 Jan 6.
86	Not critically ill patients	Jia Z.-Y., Yang J., Xia Y., Tong D.-N., Zaloga G.P., Qin H.-L., Qiu Z., Li W., Jiang X., Peng B., Xia J., Xiu D., Huang F., Zhang N., Pan H., Huang S., Zhou Y., Lin J., Ji J., Zhou Z., Li Z., Gao H., Zhang X., Du H. Safety and efficacy of an olive oil-based triple-chamber bag for parenteral nutrition: A prospective, randomized, multi-center clinical trial in China. <i>Nutrition Journal</i> . 2015; 14 (1).
87	Not RCT	Chen HS, Wang W, Zhang HD, Hong CY, Wen JM, Wu SN. Treating effects of Omega-3 fish oil fatty acids on sepsis-induced MODS. <i>Guangdong Med J</i> . 2011;32:3271-3

88	Not RCT	Gazzaniga AB, Day AT, Sankary H. The efficacy of a 20 per cent fat emulsion as a peripherally administered substrate. <i>Surg Gynecol Obstet.</i> 1985 May;160(5):387-92.
89	Not RCT	Barros KV, Cassulino AP, Schalch L, Della Valle Munhoz E, Manetta JA, Noakes PS, Miles EA, Calder PC, Flor Silveira VL. Supplemental intravenous n-3 fatty acids and n-3 fatty acid status and outcome in critically ill elderly patients in the ICU receiving enteral nutrition. <i>Clin Nutr.</i> 2013 Aug;32(4):599-605.
90	Not RCT	Barros KV, Cassulino AP, Schalch L, Munhoz ED, Manetta JA, Calder PC, Silveira VL. Pharmaconutrition: acute Fatty Acid modulation of circulating cytokines in elderly patients in the ICU. <i>JPEN J Parenter Enteral Nutr.</i> 2014 May;38(4):467-74.
91	Not RCT	Heller, A.R.; Rossler, S.; Litz, R.J.; Stehr, S.N.; Heller, S.C.; Koch, R. Omega-3 fatty acids improve the diagnosis-related clinical outcome. <i>Crit. Care. Med.</i> 2006, 34, 972–979.
92	Not RCT	Lukach VN, Girsh AO, Tolkach AB, Osipovskii NA, Iudakova TN, Glushchenko AV. [Effective use of third generation fat emulsion in total parenteral nutrition program in patients with severe sepsis] <i>Anesteziol Reanimatol.</i> 2011;(2):30-34.
93	Not RCT	Ni C, Cao J, Li D, Wu W, Cao L, Zhu C. Parenteral nutrition effects of Omega-3 fatty acids on C-reactive protein, high-density lipoprotein, lymphocyte characteristics and the treatment of critically ill patients. <i>Cell Mol Biol (Noisy-le-grand).</i> 2020;66(3):52-56. Published 2020 Jun 5.
94	Rapid vs slow lipid infusion	Suchner U, Katz DP, Fürst P, et al. Effects of intravenous fat emulsions on lung function in patients with acute respiratory distress syndrome or sepsis. <i>Crit Care Med.</i> 2001;29(8):1569-1574. doi:10.1097/00003246-200108000-00012
95	Subanalysis of Hall 2014	Hall TC, Bilku DK, Neal CP, Cooke J, Fisk HL, Calder PC, Dennison AR. The impact of an omega-3 fatty acid rich lipid emulsion on fatty acid profiles in critically ill septic patients. <i>Prostaglandins Leukot Essent Fatty Acids.</i> 2016 Sep;112:1-11.
96	Systematic Review. Individual studies screened.	van der Meij BS, van Bokhorst-de van der Schueren MA, Langius JA, Brouwer IA, van Leeuwen PA. n-3 PUFAs in cancer, surgery, and critical care: a systematic review on clinical effects, incorporation, and washout of oral or enteral compared with parenteral supplementation. <i>Am J Clin Nutr.</i> 2011 Nov;94(5):1248-65.
97	Systematic Review. Individual studies screened.	Manzanares W, Langlois PL, Dhaliwal R, Lemieux M, Heyland DK. Intravenous fish oil lipid emulsions in critically ill patients: an updated systematic review and meta-analysis. <i>Crit Care.</i> 2015;19(1):167. Published 2015 Apr 16. doi:10.1186/s13054-015-0888-7
98	Systematic Review. Individual studies screened.	Manzanares W, Dhaliwal R, Jurewitsch B, Stapleton RD, Jeejeebhoy KN, Heyland DK. Parenteral Fish Oil Lipid Emulsions in the Critically Ill: A Systematic Review and Meta-Analysis. <i>JPEN J Parenter Enteral Nutr.</i> 2013 Apr 22.
99	Systematic Review. Individual studies screened.	Manzanares W, Dhaliwal R, Jurewitsch B, Stapleton RD, Jeejeebhoy KN, Heyland DK. Alternative lipid emulsions in the critically ill: a systematic review of the evidence. <i>Intensive Care Med.</i> 2013 Jun 29.
100	Systematic Review. Individual studies screened.	Palmer AJ, Ho CK, Ajibola O, Avenell A. The role of ω -3 fatty acid supplemented parenteral nutrition in critical illness in adults: a systematic review and meta-analysis. <i>Crit Care Med.</i> 2013 Jan;41(1):307-16.
101	Systematic Review. Individual studies screened.	Zhu D, Zhang Y, Li S, Gan L, Feng H, Nie W. Enteral omega-3 fatty acid supplementation in adult patients with acute respiratory distress syndrome: a systematic review of randomized controlled trials with meta-analysis and trial sequential analysis. <i>Intensive Care Med.</i> 2014 Apr;40(4):504-12.
102	Systematic Review. Individual studies screened.	Wan X, Gao X, Bi J, Tian F, Wang X. Use of n-3 PUFAs can decrease the mortality in patients with systemic inflammatory response syndrome: a systematic review and meta-analysis. <i>Lipids Health Dis.</i> 2015 Mar 31;14:23.

103	Systematic Review. Individual studies screened.	Lu C, Sharma S, McIntyre L, Rhodes A, Evans L, Almenawer S, Leduc L, Angus DC, Alhazzani W. Omega-3 supplementation in patients with sepsis: a systematic review and meta-analysis of randomized trials. <i>Ann Intensive Care</i> . 2017 Dec;7(1):58.
104	Systematic Review. Individual studies screened.	Pradelli L, Mayer K, Klek S, et al. ω -3 Fatty-Acid Enriched Parenteral Nutrition in Hospitalized Patients: Systematic Review With Meta-Analysis and Trial Sequential Analysis. <i>JPEN J Parenter Enteral Nutr</i> . 2020;44(1):44-57. doi:10.1002/jpen.1672
105	Systematic Review. Individual studies screened.	Chen H, Wang S, Zhao Y, Luo Y, Tong H, Su L. Correlation analysis of omega-3 fatty acids and mortality of sepsis and sepsis-induced ARDS in adults: data from previous randomized controlled trials. <i>Nutr J</i> . 2018;17(1):57. Published 2018 May 31. doi:10.1186/s12937-018-0356-8
106	Systematic Review. Individual studies screened.	Chen W, Jiang H, Zhou ZY, et al. Is omega-3 fatty acids enriched nutrition support safe for critical ill patients? A systematic review and meta-analysis. <i>Nutrients</i> . 2014;6(6):2148-2164. Published 2014 May 30. doi:10.3390/nu6062148
107	Systematic Review. Individual studies screened.	Tao W, Li PS, Shen Z, Shu YS, Liu S. Effects of omega-3 fatty acid nutrition on mortality in septic patients: a meta-analysis of randomized controlled trials. <i>BMC Anesthesiol</i> . 2016;16(1):39. Published 2016 Jul 18. doi:10.1186/s12871-016-0200-7
108	Systematic Review. Individual studies screened.	Wu GH, Zaniolo O, Schuster H, Schlotzer E, Pradelli L. Structured triglycerides versus physical mixtures of medium- and long-chain triglycerides for parenteral nutrition in surgical or critically ill adult patients: Systematic review and meta-analysis. <i>Clin Nutr</i> . 2017;36(1):150-161. doi:10.1016/j.clnu.2016.01.004
109	Systematic Review. Individual studies screened.	Pradelli L, Klek S, Mayer K, et al. Omega-3 fatty acid-containing parenteral nutrition in ICU patients: systematic review with meta-analysis and cost-effectiveness analysis. <i>Crit Care</i> . 2020;24(1):634. Published 2020 Nov 3. doi:10.1186/s13054-020-03356-w
110	Too few ICU patients (only 8-14% patients ventilated)	Hajdú N, Belágyi T, Issekutz A, Bartek P, Gartner B, Oláh A. [Intravenous glutamine and early nasojejunal nutrition in severe acute pancreatitis – a prospective randomized clinical study]. <i>Magy Seb</i> . 2012 Apr;65(2):44-51. Hungarian.
111	Too few ICU patients (personal communication J Allard)	Salazar E, Alenezi S, Schwenger KJP, et al. Olive oil-based lipid emulsion is noninferior to soybean oil-based lipid emulsion in the acute care setting: A double-blind randomized controlled trial [published online ahead of print, 2021 Apr 18]. <i>Nutrition</i> . 2021;89:111283. doi:10.1016/j.nut.2021.111283
112	Unable to retrieve full-text	Liang X, Fang LZ, Zhang Y. Inhibitive effects of Omega-3 fish oil to severe sepsis. <i>China Prac Med</i> . 2009;4:158–9. [found in systematic review: Chen et al. <i>Nutrition Journal</i> (2018) 17:57]
113	PN composition varied by glucose and lipid	Huschak G, Zur Nieden K, Hoell T, Riemann D, Mast H, Stuttmann R. Olive oil based nutrition in multiple trauma patients: a pilot study. <i>Intensive Care Med</i> . 2005;31(9):1202-1208. doi:10.1007/s00134-005-2727-9